

moving objects are in view³. Heading perception is much more robust for adverse conditions such as these^{2,3}, when the subject makes real rather than simulated eye movements.

A. V. van den Berg

*Department of Physiology,
Erasmus University Rotterdam,
PO Box 1738,
3000 DR Rotterdam,
The Netherlands*

Time for tea

SIR — M. Spiro and D. Jaganyi (*Nature* **364**, 581; 1993) suggest that a chemical reaction takes place between tea and the suspension of calcium carbonate present in boiled hard water. I believe, however, that the main process taking place is physical adsorption of the coloured tannin from the tea on to the surface of the fine suspension of calcium carbonate that then settles on the surfaces of teapots and cups as a brown stain. Any chemical reaction is negligible. It is common practice for tea drinkers to add citric acid in the form of a slice of lemon to remove the calcium carbonate and prevent the formation of a stain.

I have extended the process by adding a small quantity of citric acid crystals to the hard water before boiling. The calcium bicarbonate hardness is converted into soluble citrate and no precipitate is produced after boiling.

P. P. Jones

*Cercol Laboratories,
Conways Drive,
Poole,
Dorset BH14 OPL, UK*

Velocities in the mantle

SIR — The Scientific Correspondence¹ from Wright quoted my News and Views article² out of context, and its complaints were unwarranted. My article, about the recent results of Vidale and Benz³, explained that while regions of anomalously fast seismic compressional (P) and shear (S) velocities have been found in many places at the base of the mantle with many different data sources, the fast P-velocity core-mantle boundary feature found by Vidale and Benz was in a region (beneath Alaska) that had previously revealed slow P velocities⁴⁻⁶. Material exhibiting fast P-wave velocities certainly exists at the base of the mantle, but it was not expected in this region. The statement that led to confusion, "The results are all the more intriguing because they do not occur with previous studies", was therefore meant in a regional and not a global context.

Wright interprets this sentence as an

implication that no other regions of fast seismic velocities had ever been found, says that my statement is incorrect, and cites some examples of his own work in which he found fast seismic zones at the base of the mantle in other parts of the world. His comments are unjustified, because the rest of the paragraph in my News and Views that contains the above quote lists nine other studies within the past 6 years in which fast seismic zones at the base of the mantle are reported. I am well aware, and made it clear, that the base of the mantle can display high-velocity zones. I certainly recognize the contributions that Wright has made to this subject, but point out that News and Views articles are intended to put pieces of current research into context. They are not designed to give a thorough historical background to the subject, nor could this be done with a dozen or fewer references.

Michael E. Wyession

*Department of Earth and Planetary
Sciences,
Washington University,
St Louis, Missouri 63130, USA*

1. Wright, C. *Nature* **364**, 294 (1993).
2. Wyession, M. E. *Nature* **361**, 495-496 (1993).
3. Vidale, J. E. & Benz, J. M. *Nature* **361**, 529-532 (1993).
4. Wyession, M. E. *J. geophys. Res.* **97**, 8749-8764 (1992).
5. Inoue, H., Fukao, Y., Tanabe, K. & Ogata, Y. *Phys. Earth planet. Inter.* **59**, 294-328 (1990).
6. Pulliam, R. J., Vasco, D. W. & Johnson, L. R. *J. geophys. Res.* **98**, 669-734 (1993).

OH in Saturn's rings

SIR — The presence of the hydroxyl radical (OH) near the inner satellites of Saturn is not unexpected as the adjacent moons and rings are predominantly water ice. Nevertheless, the discovery^{1,2} of OH is surprising because the amount observed implies a production rate of the H₂O parent molecule 20 times greater than theoretical calculations would suggest¹. Typical sources for circumplanetary H₂O include collisions of interplanetary micrometeoroids into inner satellites and rings, and sputtering of these objects by ions and neutral particles. To account for their observations, Shemansky *et al.*¹ suggest that the interplanetary micrometeoroid flux to the saturnian system could be 20 times greater than currently believed. But such an increased flux would dramatically shorten lifetimes for the main saturnian rings to darken, for the rings to spread by angular momentum transfer and for ring particles to erode, which would make theories of the rings' primordial origin untenable. Instead, we suggest that H₂O is primarily produced by collisions of E-ring grains into the inner satellites of Saturn.

The faint E ring is composed of

micrometre-sized icy debris, which is thought to be chips off the inner saturnian satellites^{3,4}. The orbits of these tiny grains are strongly perturbed by the non-gravitational forces of electromagnetism and radiation pressure, which cause initially circular orbits periodically to become highly elliptical³. These elongate paths bring E-ring grains across the nearly circular orbits of the inner satellites so that an average E-ring grain will strike one of the moons with a typical velocity of about 2.5 km s⁻¹ in about 20 years. These hypervelocity impacts eject relatively large amounts of vapour and sub-micrometre-sized debris into orbit around Saturn.

We now compare the relative importance of the E ring and interplanetary micrometeoroid contributions to the production of H₂O. Using values from refs 5 and 6, the total mass fluxes to the satellite Enceladus are estimated as follows:

$$\begin{aligned} \text{Interplanetary micrometeoroid flux:} \\ (\text{IP flux density}) (\text{area of Enceladus}) = \\ (4.5 \times 10^{-17} \text{ g cm}^{-2} \text{ s}^{-1}) \pi (250 \text{ km})^2 = \\ 0.088 \text{ g s}^{-1} \end{aligned}$$

$$\begin{aligned} \text{E-ring particle flux:} \\ (\text{mass of E ring}) / \\ (\text{collisional timescale with Enceladus}) = \\ (7 \times 10^{11} \text{ g}) / (20 \text{ yr}) = 1,100 \text{ g s}^{-1} \end{aligned}$$

it is seen that the flux of E-ring grains to Enceladus exceeds the nominal flux of interplanetary micrometeoroids by a factor of about 10,000. Even after accounting for the larger collision speeds of the interplanetary particles (about 25 km s⁻¹), we predict that the E-ring grains will produce collisional products at least 100 times more efficiently than their interplanetary counterparts.

Mutual collisions among E-ring particles also occur relatively frequently and at high velocity. These grain-grain collisions further enhance the efficiency at which micrometre and sub-micrometre-sized collisional products are converted into H₂O vapour. Incorporating E-ring particles into gas production calculations, we find that the amount of material blasted from the satellites is sufficient to account for the missing factor of 20 in the water production rates.

Douglas P. Hamilton

Joseph A. Burns

*Astronomy Department,
Cornell University,
Ithaca,
New York 14853, USA*

1. Shemansky, D. E., Matheson, P., Hall, D. T., Hu, H.-Y. & Tripp, T. M. *Nature* **363**, 329-331 (1993).
2. Johnson, R. E. *Nature* **363**, 300-301 (1993).
3. Horanyi, M., Burns, J. A. & Hamilton, D. P. *Icarus* **97**, 248-259 (1992).
4. Hamilton, D. P. & Burns, J. A. *Bull. Am. astr. Soc.* **24**, 1031 (1992).
5. Showalter, M. R., Cuzzi, J. N. & Larson, S. M. *Icarus* **94**, 451-473 (1991).
6. Durisen, R. H., Bode, P. W., Cuzzi, J. N., Cederbloom, S. E. & Murphy, B. W. *Icarus* **100**, 364-393 (1992).