

THE LOW CO CONTENT OF THE EXTREMELY METAL POOR GALAXY I ZW 18

ADAM LEROY¹, JOHN CANNON^{1,2}, FABIAN WALTER¹, ALBERTO BOLATTO³, AXEL WEISS⁴

Draft version February 1, 2008

ABSTRACT

We present sensitive molecular line observations of the metal-poor blue compact dwarf I Zw 18 obtained with the IRAM Plateau de Bure interferometer. These data constrain the CO $J = 1 \rightarrow 0$ luminosity within our 300 pc (FWHM) beam to be $L_{\text{CO}} < 1 \times 10^5 \text{ K km s}^{-1} \text{ pc}^2$ ($I_{\text{CO}} < 1 \text{ K km s}^{-1}$), an order of magnitude lower than previous limits. Although I Zw 18 is starbursting, it has a CO luminosity similar to or less than nearby low-mass irregulars (e.g. NGC 1569, the SMC, and NGC 6822). There is less CO in I Zw 18 relative to its B -band luminosity, H I mass, or star formation rate than in spiral or dwarf starburst galaxies (including the nearby dwarf starburst IC 10). Comparing the star formation rate to our CO upper limit reveals that unless molecular gas forms stars much more efficiently in I Zw 18 than in our own galaxy, it must have a very low CO-to-H₂ ratio, $\sim 10^{-2}$ times the Galactic value. We detect 3mm continuum emission, presumably due to thermal dust and free-free emission, towards the radio peak.

Subject headings: galaxies: individual (I Zw 18); galaxies: ISM; galaxies: dwarf, radio lines: ISM

1. INTRODUCTION

With the lowest nebular metallicity in the nearby universe ($12 + \log O/H \approx 7.2$, Skillman & Kennicutt 1993), the blue compact dwarf I Zw 18 plays an important role in our understanding of galaxy evolution. Vigorous ongoing star formation implies the presence of molecular gas, but direct evidence has been elusive. Vidal-Madjar et al. (2000) showed that there is not significant diffuse H₂, but Cannon et al. (2002) found $\sim 10^3 M_{\odot}$ of dust organized in clumps with sizes 50 – 100 pc. Vidal-Madjar et al. (2000) did not rule out compact, dense molecular clouds, and Cannon et al. (2002) argued that this dust may indicate the presence of molecular gas.

Observations by Arnault et al. (1988) and Gondhalekar et al. (1998) failed to detect CO $J = 1 \rightarrow 0$ emission, the most commonly used tracer of H₂. This is not surprising. The low dust abundance and intense radiation fields found in I Zw 18 may have a dramatic impact on the formation of H₂ and structure of molecular clouds. A large fraction of the H₂ may exist in extended envelopes surrounding relatively compact cold cores. In these envelopes, H₂ self-shields while CO is dissociated (Maloney & Black 1988). The result may be that in such galaxies [CII] or FIR emission trace H₂ better than CO (Madden et al. 1997; Israel 1997a; Pak et al. 1998). Further, H₂ may simply be underabundant, as there is a lack of grains on which to form while photodissociation is enhanced by an intense UV field. Indeed, Bell et al. (2006) found that at $Z = Z_{\odot}/100$, a molecular cloud may take as long as a Gyr to reach chemical equilibrium.

A low CO content in I Zw 18 is then expected, and a stringent upper limit would lend observational support to predictions for molecular cloud structure at low metallicity. However, while the existing upper limits are sensitive

in an absolute sense, they do not even show I Zw 18 to have a lower *normalized* CO content than a spiral galaxy (e.g. less CO per B -band luminosity). The low luminosity ($M_B \approx -14.7$, Gil de Paz et al. 2003) and large distance ($d=14$ Mpc, Izotov & Thuan 2004) of this system require very sensitive observations to set a meaningful upper limit.

In this letter we present observations, obtained with the IRAM Plateau de Bure Interferometer (PdBI)⁵, that constrain the CO luminosity, L_{CO} , to be equal to or less than that of nearby CO-poor (non-starbursting) dwarf irregulars.

2. OBSERVATIONS

I Zw 18 was observed with the IRAM Plateau de Bure Interferometer on 17, 21, and 27 April and 13 May 2004 for a total of 11 hours. The phase calibrators were 0836+710 ($F_{\nu}(115\text{GHz}) \approx 1.1 \text{ Jy}$), and 0954+556 ($F_{\nu}(115\text{GHz}) \approx 0.35 \text{ Jy}$). One or more calibrators with known fluxes were also observed during each track. The data were reduced at the IRAM facility in Grenoble using the GILDAS software package; maps were prepared using AIPS. The final CO $J = 1 \rightarrow 0$ data cube has beam size $5.59'' \times 3.42''$, and a velocity (frequency) resolution of 6.5 km s^{-1} (2.5 MHz). The velocity coverage stretches from $v_{\text{LSR}} \approx 50$ to 1450 km s^{-1} . The data have an RMS noise of $3.77 \text{ mJy beam}^{-1}$ (18 mK; $1 \text{ Jy beam}^{-1} = 4.8 \text{ K}$). The $44''$ (FWHM) primary beam completely covers the galaxy. Based on variation of the relative fluxes of the calibrators, we estimate the gain uncertainty to be $< 15\%$.

3. RESULTS

3.1. Upper Limit on CO Emission

To search for significant CO emission, we smooth the cube to 20 km s^{-1} velocity resolution, a typical line width for CO at our spatial resolution (e.g., Helfer et al. 2003).

⁵ Based on observations carried out with the IRAM Plateau de Bure Interferometer. IRAM is supported by INSU/CNRS (France), MPG (Germany) and IGN (Spain)."

¹ Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117, Heidelberg, Germany; email: leroy@mpia-hd.mpg.de

² Astronomy Department, Wesleyan University, Middletown, CT 06459, cannon@astro.wesleyan.edu

³ Radio Astronomy Lab, UC Berkeley, 601 Campbell Hall, Berkeley, CA, 94720

⁴ MPIfR, Auf dem Hügel 69, 53121, Bonn, Germany




FIG. 1.— CO $1 \rightarrow 0$ spectra of I Zw 18 towards the radio continuum/H α peak (left) and the highest significance spectra (right), which is still too faint to classify as more than marginal. The locations of both spectra are shown in Figure 2. Dashed horizontal lines show the magnitude of the RMS noise.

The noise per channel map in this smoothed cube is $\sigma_{20} \approx 0.25$ K km s $^{-1}$. Over the H I velocity range (710 – 810 km s $^{-1}$, van Zee et al. 1998), there are no regions with $I_{\text{CO},20} > 1$ K km s $^{-1}$ (4σ) within the primary beam. We pick a slightly conservative upper limit for two reasons. First, if there were CO emission with this intensity we would be certain of detecting it. Second, the noise in the cube is slightly non-Gaussian, so that the false positive rate for $I_{\text{CO},20} > 1$ K km s $^{-1}$ — estimated from the negatives and the channel maps outside the H I velocity range — is $\sim 0.2\%$, very close to that of a 3σ deviate.

For $d = 14$ Mpc, the synthesized beam has a FWHM of 300 pc and an area of 1.0×10^5 pc 2 . Our intensity limit, $I_{\text{CO}} < 1$ K km s $^{-1}$, therefore translates to a CO luminosity limit of $L_{\text{CO}} < 1 \times 10^5$ K km s $^{-1}$ pc 2 .

There is a marginal signal toward the southern knot of H α emission ($9^{\text{h}}34^{\text{m}}02^{\text{s}}.4$, $55^{\circ}14'23''.0$). This emission has the largest $|I_{\text{CO},20}|$ found over the H I velocity range, corresponding to $L_{\text{CO}} \sim 8 \times 10^4$ K km s $^{-1}$ pc 2 , just below our limit. This same line of sight also shows $|I_{\text{CO}}| > 2\sigma$ over three consecutive channels, a feature seen along only one other line of sight (in negative) over the H I velocity range. The marginal signal is suggestively located in the southeast of I Zw 18, where Cannon et al. (2002) identified several potential sites of molecular gas from regions of relatively high extinction. While tantalizing, the signal is not strong enough to be categorized as a detection. Figure 1 shows CO spectra towards the H α /radio continuum peak (Cannon et al. 2002, 2005; Hunt et al. 2005a, see Figure 2) and this marginal signal.

3.2. Continuum Emission

We average the data over all channels and produce a continuum map with noise $\sigma_{115\text{GHz}} = 0.35$ mJy beam $^{-1}$. The highest value in the map is $I_{115\text{GHz}} = 1.06 \pm 0.35$ mJy beam $^{-1}$ at $\alpha_{2000} = 9^{\text{h}}34^{\text{m}}02^{\text{s}}.1$, $\delta_{2000} = +55^{\circ}14'27''.0$. This is within a fraction of a beam of the 1.4 GHz peak identified by Cannon et al. (2005), $\alpha_{2000} = 9^{\text{h}}34^{\text{m}}02^{\text{s}}.1$, $\delta_{2000} = +55^{\circ}14'28''.06$ and Hunt et al. (2005a, $\alpha_{2000} = 9^{\text{h}}34^{\text{m}}02^{\text{s}}$, $\delta_{2000} = +55^{\circ}14'29''.06$). Figure 2 shows the radio continuum peak and 115 GHz continuum contours plotted over H α emission from I Zw 18 (Cannon et al. 2002). There is only one other region with $|I_{115\text{GHz}}| > 3\sigma_{115\text{GHz}}$ within the primary beam and the star-forming extent of I Zw 18 occupies $\approx 10\%$ of the primary beam. Therefore, we estimate the chance of a false positive coincident with the galaxy to be only $\sim 10\%$.

4. DISCUSSION

Here we discuss the implications of our CO upper limit and continuum detection. We adopt the following properties for I Zw 18, all scaled to $d = 14$ Mpc: $M_B = -14.7$ (Gil de Paz et al. 2003), $M_{\text{HI}} = 1.4 \times 10^8$ M $_{\odot}$ (van Zee et al. 1998), H α luminosity $\log_{10} H_{\alpha} = 39.9$ erg s $^{-1}$ (Cannon et al. 2002; Gil de Paz et al. 2003), 1.4 GHz flux $F_{1.4} = 1.79$ mJy (Cannon et al. 2005).

4.1. Point Source Luminosity

Our upper limit along each line of sight, $L_{\text{CO}} < 1 \times 10^5$ K km s $^{-1}$ pc 2 , matches the luminosity of a fairly massive Galactic giant molecular cloud (Blitz 1993). For a Galactic CO-to-H $_2$ conversion factor, 2×10^{20} cm $^{-2}$ (K km s $^{-1}$) $^{-1}$, the corresponding molecular gas mass is $M_{\text{Mol}} \approx 4.4 \times 10^5$ M $_{\odot}$, similar to the mass of the Orion-Monoceros complex (e.g. Wilson et al. 2005).

4.2. Comparison With More Luminous Galaxies

In galaxies detected by CO surveys, the CO content per unit B -band luminosity is fairly constant. Figure 3 shows the CO luminosity normalized by B -band luminosity, L_{CO}/L_B , as a function of absolute B -band magnitude (L_B is extinction corrected). L_{CO}/L_B is nearly constant over two orders of magnitude in L_B , though with substantial scatter (much of it due to the extrapolation from a single pointing to L_{CO}).

Based on these data and assuming that L_{CO} is not a function of the metallicity of the galaxy, we may extrapolate to an expected CO luminosity for I Zw 18. For $M_{B,\text{IZw18}} \approx -14.7$ the CO luminosity corresponding to the median value of L_{CO}/L_B (dashed line) in Figure 3 is $L_{\text{CO,IZw18}} \approx 1.7 \times 10^6$ K km s $^{-1}$ pc 2 . The H α , 1.4 GHz, and H I luminosities lead to similar predictions. Young et al. (1996) found $M_{\text{H}_2}/L_{\text{H}\alpha} \approx 10 L_{\odot}/M_{\odot}$ for Sd-Irr galaxies, which implies $L_{\text{CO,IZw18}} \sim 4 \times 10^6$ K km s $^{-1}$ pc 2 . Murgia et al. (2005) measured $F_{\text{CO}}/F_{1.4} \approx 10$ Jy km s $^{-1}$ (mJy) $^{-1}$ for spirals, that would imply $L_{\text{CO,IZw18}} \sim 10^7$ K km s $^{-1}$. For Sd/Sm galaxies, $M_{\text{H}_2}/M_{\text{HI}} \approx 0.2$ (Young & Scoville 1991), leading to $L_{\text{CO,IZw18}} \sim 5 \times 10^6$ K km s $^{-1}$ pc 2 . Both $M_{\text{H}_2}/L_{\text{H}\alpha}$ and $M_{\text{H}_2}/M_{\text{HI}}$ tend to be even higher in earlier-type spirals.

Therefore, surveys would predict $L_{\text{CO,IZw18}} \gtrsim 2 \times 10^6$ K km s $^{-1}$ pc 2 , very close to the previously established upper limits of $2 - 3 \times 10^6$ K km s $^{-1}$ pc 2 (Arnault et al. 1988; Gondhalekar et al. 1998). With the present observations, we constrain $L_{\text{CO}} < 1 \times 10^5$ K km s $^{-1}$ pc 2 and thus clearly rule out $L_{\text{CO}} \sim 10^6$ K km s $^{-1}$ pc 2 . This may be seen in Figure 3; even if I Zw 18 has the highest possible CO content, it will still have a lower L_{CO}/L_B than 97% of the survey galaxies.

4.3. Comparison With Nearby Metal-Poor Dwarfs

The subset of irregular galaxies detected by CO surveys tend to be CO-rich and actively star-forming, resembling scaled-down versions of spiral galaxies (Young et al. 1995, 1996; Leroy et al. 2005). Such galaxies may not be representative of all dwarfs. Because they are nearby, several of the closest dwarf irregulars have been detected despite very small L_{CO} . With their low masses and metallicities, they may represent good points of comparison for I Zw 18. Table 1 and Figure 3 show CO lumi-




FIG. 2.— V-band and H α (right, Cannon et al. 2002) images of I Zw 18. Overlays on the left image show the size of the synthesized beam and the locations of the spectra shown in Figure 1. Contours on the right image show continuum emission in increments of 0.5σ significance and the location of the radio continuum peak. The primary beam is larger than the area shown. Both optical maps are on linear stretches. V-band data obtained from the MAST Archive, originally observed for GO program 9400, PI: T. Thuan).

nosities and L_{CO}/L_B for four nearby dwarfs: NGC 1569, the Small Magellanic Cloud (SMC), NGC 6822, and IC 10. The SMC, NGC 1569, and NGC 6822 have $L_{\text{CO}} \sim 10^5 \text{ K km s}^{-1} \text{ pc}^2$, close to our upper limit, and occupy a region of L_{CO}/L_B - L_B parameter space similar to I Zw 18. All four of these galaxies have active star formation but very low CO content relative to their other properties.

We test whether our observations would have detected CO in NGC 1569, the SMC, and IC 10 at the plausible lower limit of 10 Mpc (from $H_0 = 72 \text{ km s}^{-1}$) or our adopted distance of 14 Mpc. We convolve the integrated intensity maps to resolutions of 210 and 300 pc and measure the peak integrated intensity. The results appear in columns 4 and 5 of Table 1. The PdBI observations of NGC 1569 resolve out most of the flux, so we also apply this test to a distribution with the size and luminosity derived by Greve et al. (1996) from single dish observations. Our observations would detect an analog to IC 10 but not the SMC, with NGC 1569 an intermediate case. With a factor of ~ 3 better sensitivity (requiring ~ 10 times more observing time) we would expect to detect all three nearby galaxies. However, achieving such sensitivity with present instrumentation will be quite challenging. ALMA will likely be necessary to place stronger constraints on CO in galaxies like I Zw 18.

IC 10 may be the nearest blue compact dwarf (Richer et al. 2001), so it may be telling that we would detect it at the distance of I Zw 18. The blue compact galaxies that have been detected in CO have L_{CO}/L_B similar to IC 10 (Gondhalekar et al. 1998, the diamonds in Figure 3). Most searches for CO towards BCDs have yielded nondetections, so those detected may not be representative, but I Zw 18 is clearly not among the “CO-rich” portion of the BCD population.

4.4. Interpretation of the Continuum

We measure continuum intensity of $F_{115\text{GHz}} = 1.06 \pm 0.35 \text{ mJy}$ towards the radio continuum peak. The continuum is detected along only one line of sight, so we refer to it here as a point source and compare it to integrated values for I Zw 18. $F_{115\text{GHz}}$ is expected to be the product of mainly two types of emission: thermal free-free emission and thermal dust emission. At long wavelengths, the integrated thermal free-free emission is $F_{1.4\text{GHz}}(\text{free} - \text{free}) \approx 0.52 - 0.75 \text{ mJy}$ (Cannon et al. 2005; Hunt et al. 2005a), implying $F_{115\text{GHz}}(\text{free} - \text{free}) = 0.36 - 0.51 \text{ mJy}$ at 115 GHz ($F_\nu \propto \nu^{-0.1}$). The H α flux predicts a similar value, $F_{115\text{GHz}}(\text{free} - \text{free}) = 0.34 \text{ mJy}$ (Cannon et al. 2005, Equation 1). Hunt et al. (2005b) placed an upper limit of $F_\nu(850) < 2.5 \text{ mJy}$ on dust continuum emission at $850\mu\text{m}$; this is consistent with the $\sim 5 \times 10^3 M_\odot$ estimated by Cannon et al. (2002) given almost any reasonable dust properties. Extrapolating this to 2.6 mm assuming a pure blackbody spectrum, the shallowest plausible SED, constrains thermal emission from dust to be $< 0.25 \text{ mJy}$ at 115 GHz. Based on these data, we would predict $F_{115\text{GHz}} \lesssim 0.75 \text{ mJy}$. Thus our measured $F_{115\text{GHz}}$ is consistent with, but somewhat higher than, the thermal free-free plus dust emission expected based on optical, centimeter, and submillimeter data.

4.5. Relation to Star Formation

I Zw 18 has a star formation rate $\sim 0.06 - 0.1 M_\odot \text{ yr}^{-1}$, based on H α and cm radio continuum measurements (Cannon et al. 2002; Kennicutt 1998a; Hunt et al. 2005a). Our continuum flux suggests a slightly higher value $\approx 0.15 - 0.2 M_\odot \text{ yr}^{-1}$ (following Hunt et al. 2005a; Condon 1992), with the exact value depending on the contribution from thermal dust emission. For any value in this range, the star formation rate per CO luminosity, SFR/L_{CO} is much higher in I Zw 18 than in spirals. For comparison, our upper limit and the molecular “Schmidt Law” derived by Murgia et al. (2002) predicts a star for-




FIG. 3.— CO luminosity normalized by absolute blue magnitude for galaxies with Hubble Type Sb or later (black circles, Young et al. 1995; Elfhag et al. 1996; Böker et al. 2003; Leroy et al. 2005). We also plot nearby dwarfs from Table 1 (crosses) and blue compact galaxies compiled by Gondhalekar et al. (1998, , diamonds). The shaded regions shows our upper limit for I Zw 18, with the range in M_B for distances from 10 to 20 Mpc. The dashed line and light shaded region show the median value and 1σ scatter in L_{CO}/L_B for spirals and dwarf starbursts. *Methodology:* We extrapolate from I_{CO} in central pointings to L_{CO} assuming the CO to have an exponential profile with scale length $0.1 d_{25}$ (Young et al. 1995), including only galaxies where the central pointing measures $> 20\%$ of L_{CO} . We adopt B magnitudes (corrected for internal and Galactic extinction), distances (Tully-Fisher when available, otherwise Virgocentric-flow corrected Hubble flow), and radii from LEDA (Paturel et al. 2003).

mation rate $\lesssim 2 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$. Fits by Young et al. (1996) and Kennicutt (1998b, applied to just the molecular limit) yield similar values. Again, I Zw 18 is similar to the SMC and NGC 6822, which have star formation rates of $0.05 M_{\odot} \text{ yr}^{-1}$ and $0.04 M_{\odot} \text{ yr}^{-1}$ (Wilke et al. 2004; Israel 1997b) and $L_{CO} \sim 10^5 \text{ K km s}^{-1} \text{ pc}^2$.

4.6. Variations in X_{CO}

Several calibrations of the CO-to- H_2 conversion factor, X_{CO} as a function of metallicity exist in the literature. The topic has been controversial and these calibrations range from little or no dependence (e.g. Walter 2003; Rosolowsky et al. 2003) to very steep dependence (e.g., $X_{CO} \propto Z^{-2.7}$ Israel 1997a). Comparing the star formation rate to our CO upper limit, we may rule out

that I Zw 18 has a Galactic X_{CO} unless molecular gas in I Zw 18 forms stars much more efficiently than in the Galaxy. Either the ratio of CO-to- H_2 is low in I Zw 18 or molecular gas in this galaxy forms stars with an efficiency two orders of magnitude higher than that in spiral galaxies.

5. CONCLUSIONS

We present new, sensitive observations of the metal-poor dwarf galaxy I Zw 18 at 3 mm using the Plateau de Bure Interferometer. These data constrain the integrated CO $J = 1 \rightarrow 0$ intensity to be $I_{CO} < 1 \text{ K km s}^{-1}$ over our 300 pc (FWHM) beam and the luminosity to be $L_{CO} < 1 \times 10^5 \text{ K km s}^{-1} \text{ pc}^2$.

I Zw 18 has less CO relative to its B -band luminosity, H I mass, or SFR than spiral galaxies or dwarf starbursts, including more metal-rich blue compact galaxies such as IC 10 ($Z_{IC 10} \sim Z_{\odot}/4$, Lee et al. 2003). Because of its small size and large distance, these are the first observations to impose this constraint.

We show that I Zw 18 should be grouped with several local analogs — NGC 1569, the SMC, NGC 6822 — as a galaxy with active star formation but a very low CO content relative to its other properties. In these galaxies, observations suggest that the environment affects the molecular gas and these data suggest that the same is true in I Zw 18. A simple comparison of star formation rate to CO content shows that this must be true at a basic level: either the ratio of CO to H_2 is dramatically low in I Zw 18 or molecular gas in this galaxy forms stars with an efficiency two orders of magnitude higher than that in spiral galaxies.

We detect 3mm continuum with $F_{115 \text{ GHz}} = 1.06 \pm 0.35 \text{ mJy}$ coincident with the radio peak identified by Cannon et al. (2005) and Hunt et al. (2005a). This flux is consistent with but somewhat higher than the thermal free-free plus dust emission one would predict based on centimeter, submillimeter, and optical measurements.

Finally, we note that improving on this limit with current instrumentation will be quite challenging. The order of magnitude increase in sensitivity from ALMA will be needed to place stronger constraints on CO in galaxies like I Zw 18.

We thank Roberto Neri for his help reducing the data. We acknowledge the usage of the HyperLeda database (<http://leda.univ-lyon1.fr>).

REFERENCES

- Arnault, P., Kunth, D., Casoli, F., & Combes, F. 1988, *A&A*, 205, 41
- Bell, T. A., Roueff, E., Viti, S., & Williams, D. A. 2006, *MNRAS*, 371, 1865
- Blitz, L. 1993, *Protostars and Planets III*, 125
- Böker, T., Lisenfeld, U., & Schinnerer, E. 2003, *A&A*, 406, 87
- Cannon, J. M., Skillman, E. D., Garnett, D. R., & Dufour, R. J. 2002, *ApJ*, 565, 931
- Cannon, J. M., Walter, F., Skillman, E. D., & van Zee, L. 2005, *ApJ*, 621, L21
- Condon, J. J. 1992, *ARA&A*, 30, 575
- Gil de Paz, A., Madore, B. F., & Pevunova, O. 2003, *ApJS*, 147, 29
- Elfhag, T., Booth, R. S., Hoeglund, B., Johansson, L. E. B., & Sandqvist, A. 1996, *A&AS*, 115, 439
- Gondhalekar, P. M., Johansson, L. E. B., Brosch, N., Glass, I. S., & Brinks, E. 1998, *A&A*, 335, 152
- Greve, A., Becker, R., Johansson, L. E. B., & McKeith, C. D. 1996, *A&A*, 312, 391
- Helfer, T. T., Thornley, M. D., Regan, M. W., Wong, T., Sheth, K., Vogel, S. N., Blitz, L., & Bock, D. C.-J. 2003, *ApJS*, 145, 259
- Hunt, L. K., Dyer, K. K., & Thuan, T. X. 2005a, *A&A*, 436, 837
- Hunt, L., Bianchi, S., & Maiolino, R. 2005b, *A&A*, 434, 849
- Israel, F. P. 1997, *A&A*, 328, 471
- Israel, F. P. 1997, *A&A*, 317, 65
- Izotov, Y. I., & Thuan, T. X. 2004, *ApJ*, 616, 768
- Kennicutt, R. C., Jr. 1998a, *ARA&A*, 36, 189
- Kennicutt, R. C., Jr. 1998b, *ApJ*, 498, 541
- Lee, H., McCall, M. L., & Richer, M. G. 2003, *AJ*, 125, 2975

TABLE 1
CO IN NEARBY LOW MASS GALAXIES

Galaxy	M_B (mag)	L_{CO} (K km s ⁻¹ pc ²)	$I_{CO,210}^a$ (K km s ⁻¹)	$I_{CO,300}^a$ (K km s ⁻¹)	Reference
NGC 1569	-16.5	1.2×10^5	1.1	0.8	Greve et al. (1996)
	-16.5	0.2×10^5	0.8	0.5	Taylor et al. (1999)
SMC	-16	1.5×10^5	0.5	0.4	Mizuno et al. (2001, 2006)
NGC 6822	-16	1.2×10^5	Israel (1997b)
IC 10	-16.5	2.2×10^6	3.8	2.2	Leroy et al. (2006)
I Zw 18	-14.7	$< 2 \times 10^6$	Arnault et al. (1988); Gondhalekar et al. (1998)
I Zw 18	-14.7	$\lesssim 1 \times 10^5$	< 1	< 1	this paper

^a Peak integrated intensity at 210 and 300 pc, corresponding to our beam size at 10 and 14 Mpc, respectively.

Leroy, A., Bolatto, A. D., Simon, J. D., & Blitz, L. 2005, ApJ, 625, 763
 Leroy, A., Bolatto, A., Walter, F., & Blitz, L. 2006, ApJ, 643, 825
 Madden, S. C., Poglitsch, A., Geis, N., Stacey, G. J., & Townes, C. H. 1997, ApJ, 483, 200
 Maloney, P., & Black, J. H. 1988, ApJ, 325, 389
 Mizuno, N., Rubio, M., Mizuno, A., Yamaguchi, R., Onishi, T., & Fukui, Y. 2001, PASJ, 53, L45
 Mizuno, N., et al. 2006, in prep.
 Murgia, M., Crapsi, A., Moscadelli, L., & Gregorini, L. 2002, A&A, 385, 412
 Murgia, M., Helfer, T. T., Ekers, R., Blitz, L., Moscadelli, L., Wong, T., & Paladino, R. 2005, A&A, 437, 389
 Pak, S., Jaffe, D. T., van Dishoeck, E. F., Johansson, L. E. B., & Booth, R. S. 1998, ApJ, 498, 735
 Paturel, G., Petit, C., Prugniel, P., Theureau, G., Rousseau, J., Brouty, M., Dubois, P., & Cambr sy, L. 2003, A&A, 412, 45
 Richer, M. G., et al. 2001, A&A, 370, 34
 Rosolowsky, E., Engargiola, G., Plambeck, R., & Blitz, L. 2003, ApJ, 599, 258

Skillman, E. D., & Kennicutt, R. C., Jr. 1993, ApJ, 411, 655
 Taylor, C. L., Kobulnicky, H. A., & Skillman, E. D. 1998, AJ, 116, 2746
 Taylor, C. L., H ttmeister, S., Klein, U., & Greve, A. 1999, A&A, 349, 424
 van Zee, L., Westpfahl, D., Haynes, M. P., & Salzer, J. J. 1998, AJ, 115, 1000
 Vidal-Madjar, A., et al. 2000, ApJ, 538, L77
 Walter, F. 2003, IAU Symposium, 221, 176P
 Wilke, K., Klaas, U., Lemke, D., Mattila, K., Stickel, M., & Haas, M. 2004, A&A, 414, 69
 Wilson, B. A., Dame, T. M., Mashed, M. R. W., & Thaddeus, P. 2005, A&A, 430, 523
 Young, J. S., et al. 1995, ApJS, 98, 219
 Young, J. S., & Scoville, N. Z. 1991, ARA&A, 29, 581
 Young, J. S., Allen, L., Kenney, J. D. P., Lesser, A., & Rownd, B. 1996, AJ, 112, 1903