



Contents lists available at ScienceDirect

Icarus

journal homepage: [www.elsevier.com/locate/icarus](http://www.elsevier.com/locate/icarus)



## Galileo in-situ dust measurements in Jupiter's gossamer rings

Harald Krüger<sup>a,b,\*</sup>, Douglas P. Hamilton<sup>c</sup>, Richard Moissl<sup>a</sup>, Eberhard Grün<sup>b,d</sup>

<sup>a</sup> Max-Planck-Institut für Sonnensystemforschung, Max-Planck-Str. 2, 37191 Katlenburg-Lindau, Germany

<sup>b</sup> Max-Planck-Institut für Kernphysik, Postfach 103980, 69029 Heidelberg, Germany

<sup>c</sup> Astronomy Department, University of Maryland, College Park, MD 20742-2421, USA

<sup>d</sup> Laboratory for Atmospheric and Space Physics, University of Colorado, Boulder, CO 80303-7814, USA

### ARTICLE INFO

#### Article history:

Received 13 March 2008  
Revised 16 December 2008  
Accepted 23 March 2009  
Available online xxx

#### Keywords:

Dust  
Jupiter, satellites  
Jupiter, rings  
Planetary, rings

### ABSTRACT

Galileo was the first artificial satellite to orbit Jupiter. During its late orbital mission the spacecraft made two passages through the giant planet's gossamer ring system. The impact-ionization dust detector on board successfully recorded dust impacts during both ring passages and provided the first in-situ measurements from a dusty planetary ring. During the first passage—on 5 November 2002 while Galileo was approaching Jupiter—dust measurements were collected until a spacecraft anomaly at 2.33R<sub>J</sub> (Jupiter radii) just 16 min after a close flyby of Amalthea put the spacecraft into a safing mode. The second ring passage on 21 September 2003 provided ring dust measurements down to about 2.5R<sub>J</sub> and the Galileo spacecraft was destroyed shortly thereafter in a planned impact with Jupiter. In all, a few thousand dust impacts were counted with the instrument accumulators during both ring passages, but only a total of 110 complete data sets of dust impacts were transmitted to Earth. Detected particle sizes range from about 0.2 to 5 μm, extending the known size distribution by an order of magnitude towards smaller particles than previously derived from optical imaging [Showalter, M.R., de Pater, I., Verbanac, G., Hamilton, D.P., Burns, J.A., 2008. *Icarus* 195, 361–377; de Pater, I., Showalter, M.R., Macintosh, B., 2008. *Icarus* 195, 348–360]. The grain size distribution increases towards smaller particles and shows an excess of these tiny motes in the Amalthea gossamer ring compared to the Thebe ring. The size distribution for the Amalthea ring derived from our in-situ measurements for the small grains agrees very well with the one obtained from images for large grains. Our analysis shows that particles contributing most to the optical cross-section are about 5 μm in radius, in agreement with imaging results. The measurements indicate a large drop in particle flux immediately interior to Thebe's orbit and some detected particles seem to be on highly-tilted orbits with inclinations up to 20°. Finally, the faint Thebe ring extension was detected out to at least 5R<sub>J</sub>, indicating that grains attain higher eccentricities than previously thought. The drop interior to Thebe, the excess of submicron grains at Amalthea, and the faint ring extension indicate that grain dynamics is strongly influenced by electromagnetic forces. These findings can all be explained by a shadow resonance as detailed by Hamilton and Krüger [Hamilton, D.P., Krüger, H., 2008. *Nature* 453, 72–75].

© 2009 Published by Elsevier Inc.

### 1. Previous imaging results

All four giant planets of our Solar System are surrounded by huge tenuous ring systems which contain mostly micrometer- and submicrometer-sized dust particles (Burns et al., 2001). In these rings, dust densities are so low that particle collisions are negligible, and grain dynamics is substantially perturbed by non-gravitational forces. The 'dusty' rings are interesting and valuable counterpoints to the collisionally dominated opaque and dense rings of Saturn and Uranus which are populated primarily by macroscopic centimeter- to meter-sized objects.

Jupiter's ring system was investigated with remote imaging from the Earth and from the Voyager, Galileo, Cassini and New Horizons spacecraft, revealing significant structure in the ring: at least four components have been identified (Ockert-Bell et al., 1999; Burns et al., 1999; de Pater et al., 1999): the main ring, interior halo and two gossamer rings. The small moons Metis, Adrastea, Amalthea and Thebe are embedded in the ring system and act as sources of ring dust via meteoroid impact erosion of their surfaces (Burns et al., 1999). The faint gossamer rings appear to extend primarily inward from the orbits of Amalthea and Thebe (Figs. 1 and 2). In addition, the vertical limits of each moon's slightly inclined orbit very closely match the vertical extensions of these two rings (Ockert-Bell et al., 1999). These observations imply a close relationship between the rings and embedded moonlets. Outside the orbit of Thebe, a swath of faint material is seen out to about 3.75R<sub>J</sub> (Jupiter radius, R<sub>J</sub> = 71,492 km) distance from the planet.

\* Corresponding author. Address: Max-Planck-Institut für Sonnensystemforschung, Max-Planck-Str. 2, 37191 Katlenburg-Lindau, Germany.  
E-mail address: [krueger@mps.mpg.de](mailto:krueger@mps.mpg.de) (H. Krüger).

Beyond this distance, the rings fade slowly into the background. Normal optical depths are about  $10^{-6}$  for the main ring and halo, and about 10–100 times less for the Amalthea ring and Thebe rings. Analysis of the few gossamer ring images implies particle radii of 5–10  $\mu\text{m}$  with additional contributions from larger material (Showalter et al., 2008; de Pater et al., 2008). In this paper, we show that smaller grains are also present in large numbers.

The simplest picture of particle dynamics in the ring implies that dust grains ejected from the surfaces of each moon would rapidly disperse in longitude and nodal angles while maintaining their initial inclinations (Burns et al., 1999). As such material evolves inward under Poynting–Robertson drag, it would naturally produce the two overlapping rings with rectangular profiles. Support for this interpretation comes from the fact that both gossamer rings show concentrations at the vertical extremes, where particles on inclined orbits spend most of their time. The extension of Thebe's gossamer ring beyond Thebe's orbit and recently identified radial structure in the gossamer rings, however, violates this simple and elegant picture and has been attributed to an electromagnetic process involving Jupiter's intense magnetic field by Hamilton and Krüger (2008). Similarly, recent Keck imaging by de Pater et al. (2008), which highlights large backscattering ring particles, showed that the Amalthea and Thebe rings appear to be confined to regions just interior to their bounding satellites.

## 2. Galileo in-situ dust measurements

The Galileo spacecraft was the first artificial satellite of Jupiter, circling the giant planet between 1996 and 2003. Near the end of the mission, the spacecraft passed directly through the rings twice, on 5 November 2002 and 21 September 2003, offering a unique opportunity for in-situ studies of planetary rings. The in-situ dust detector on board (Grün et al., 1992) counted several thousand dust impacts during both ring passages, and the full data sets, consisting of impact direction, charge amplitudes, rise times, etc., for 110 separate impacts were transmitted to Earth. The first ring passage included a close flyby at Amalthea with a closest approach distance of 244 km, just outside the Hill sphere of this jovian moon. The flyby provided an improved mass estimate for the satellite, with an implied density of  $\sim 800 \text{ kg m}^{-3}$  (Anderson et al., 2005).

Galileo's traversal of Jupiter's gossamer rings provided the first in-situ measurements of a dusty planetary ring. In-situ dust measurements nicely complement imaging, providing important additional information about the physical properties of the dust environment. In particular, in-situ measurements constrain dust spatial densities along the spacecraft trajectory as well as grain masses, size distributions, impact speeds and grain dynamics.

In this paper we present and analyse the complete in-situ dust measurements obtained during both Galileo gossamer ring passages. We analyse grain impact directions and impact rates and derive dust number densities and grain size distributions from the measurements. We interpret results in terms of the gossamer rings' structure and the dynamics of charged ring particles.

### 2.1. Dust detection geometry

Galileo was a dual spinning spacecraft with an antenna that pointed antiparallel to the positive spin axis. The antenna usually pointed towards Earth. The Dust Detector System (DDS) was mounted on the spinning section of Galileo underneath the magnetometer boom (Kivelson et al., 1992), with the sensor axis offset by  $60^\circ$  from the positive spin axis (Krüger et al., 1999b). Fig. 3 shows a schematic view of the Galileo spacecraft and the geometry of dust detection.

The rotation angle,  $\Theta$ , measured the viewing direction of the dust sensor at the time of a dust impact. During one spin revolution

of the spacecraft,  $\Theta$  scanned through a complete circle of  $360^\circ$ . At  $\Theta \simeq 90^\circ$  and  $\simeq 270^\circ$  the sensor axis lay nearly in the ecliptic plane, and at  $0^\circ$  it was close to the ecliptic north direction. Rotation angles are taken positive around the negative spin axis of the spacecraft which pointed towards Earth. This is done to facilitate comparison of the Galileo spin angle data with those taken by Ulysses, which, unlike Galileo, had its positive spin axis pointed towards Earth (Grün et al., 1995).

The field-of-view (FOV) of the dust sensor target was  $140^\circ$ . Due to the offset of  $60^\circ$  between the sensor axis and the spacecraft spin axis, over one spacecraft spin revolution, the sensor axis scanned the surface of a cone with  $120^\circ$  opening angle centered on the anti-Earth direction. Dust particles that arrived from within  $10^\circ$  of the positive spin axis (anti-Earth direction) could be detected at all rotation angles  $\Theta$ , whereas those that arrived with angles between  $10^\circ$  and  $130^\circ$  from the positive spin axis could be detected over only a limited range of rotation angles. In the frame fixed to the spacecraft, we define the *impact angle* between the impact velocity and the sensor axis as  $\phi$ , and the angle between the impact velocity and the spacecraft's anti-Earth spin axis as  $\psi$ .

Fig. 3 shows that the magnetometer boom ([MAG]; Kivelson et al., 1992) was in the field of view of the dust sensor. The Energetic Particles Detector (EPD; Williams et al., 1992) and the Plasma Instrument (PLS; Frank et al., 1992) partially obscured the FOV of the dust sensor as well (Fig. 4). In other words, at certain spacecraft rotation angles  $\Theta$ , particles approaching at angles with respect to the spacecraft spin axis  $\psi \geq 90^\circ$  hit the boom and these Galileo instruments instead of the sensor target. The effect of this obscuration was first recognized in measurements of the jovian dust stream particles (Krüger et al., 1999b).

### 2.2. Dust impact and noise identification

Dust grains hitting the sensor target generate a plasma cloud of evaporating grain and target material. For each impact, three independent measurements of the resulting plasma cloud were used to derive the impact speed  $v$  and the mass  $m$  of the particle: the electron signal, an ion signal, and a channeltron signal (Grün et al., 1992). The charge  $Q$  released upon impact onto the target is roughly described by the relation (Göller and Grün, 1989)

$$Q \propto m \cdot v^{3.5}. \quad (1)$$

The dust instrument was empirically calibrated in the speed range  $2\text{--}70 \text{ km s}^{-1}$ . Furthermore, the coincidence times of the three charge signals together with the charges themselves are used to sort each impact into one of four classes. Class 3 impacts have three charge signals, two are required for class 2 and class 1 events, and only one for class 0 (Baguhl, 1993; Grün et al., 1995; Krüger et al., 1999a). In addition to the four classes, the dust data were categorized into six amplitude ranges of the impact-generated ion charge, each range covering one order of magnitude in charge (here denoted by AR1 to AR6; Grün et al., 1995). Hence, taking the classes and amplitude ranges together, the dust data were grouped into  $4 \times 6 = 24$  categories.

Class 3 signals, our highest quality, are real dust impacts while class 0 events are mostly noise. Class 1 and class 2 events were true dust impacts in interplanetary space (Baguhl et al., 1993; Krüger et al., 1999a). However, during Galileo's entire Jupiter mission from 1996 to 2002—while the spacecraft was in the inner jovian magnetosphere—energetic particles from the jovian plasma environment caused enhanced noise rates in class 2 and the lower quality classes. By analysing the properties of the jovian stream particles and comparing them with the noise events, the noise could be eliminated from the class 2 data (Krüger et al., 1999b, 2005). In particular, most class 0 and class 1 events detected in the jovian environment are probably noise.

Before the two ring flybys that are the subject of this paper, Galileo had only once been within  $6R_J$  of the planet, on approach in December 1995. Due to uncertainty about the effects of Jupiter's harsh radiation environment, the dust instrument was switched to a less sensitive mode to protect it (Grün et al., 1996). Accordingly, a very low noise rate was measured. The instrument's sensitivity was later increased, and for the duration of the mission, it recorded an increasing noise level with decreasing distance to the planet.

We have tested the applicability of the noise identification scheme, described in detail by Krüger et al. (1999b) and Krüger et al. (2005), to the near-Jupiter region and improved upon it. A modified noise identification scheme was derived for the gossamer ring data by Moissl (2005), showing that class 1 also contains likely candidates for real dust impacts. For class 2, AR1 only the target-ion grid coincidence was used as a criterion for noise events (i.e.  $EIC = 0$ ) while for the higher amplitude ranges (AR2–6) the scheme of Krüger et al. (2005) was applied unchanged (i.e.  $[EA-IA \leq 1$  or  $EA-IA \geq 7]$  and  $CA \leq 2$ ; EA, IA and CA are the digital values of the charge amplitudes measured on the target, ion grid and channeltron, respectively—see Grün et al., 1995 for a description of these parameters). For class 1 the following criterion for noise events was used independent of the amplitude range of the event:  $[EA-IA \leq 2$  or  $EA-IA \geq 9]$  and  $CA \leq 2$ . More details of the noise identification in the gossamer ring data will be described by Krüger et al. (in preparation).

We use this scheme throughout this paper to separate noise events from true dust impacts. Note that this noise removal technique uses statistical arguments and is applicable to large data sets only; individual dust impacts may be erroneously classified as noise and vice versa.

### 2.3. Instrument operation and data transmission

Galileo had a very low data transmission capability because of the failure of its high-gain antenna to open completely. For the dust measurements this meant that the full set of parameters measured during a dust particle impact or noise spike could only be transmitted to Earth for a limited number of events. The data sets of all other events (whether noise or true impacts) were lost. All events (dust and noise), however, were always counted with one of the 24 accumulators (Grün et al., 1995) as described in Section 2.2. This allows us to correct the dust measurements for incomplete data transmission and to derive reliable event rates. In particular, no indications for unrecognized accumulator overflows were seen in the data from both gossamer ring passages as has been problematic for some other stages of the mission (Krüger et al., 2001).

Galileo dust data could be read out from the instrument memory with different rates (see Krüger et al. (2001), for a description). In order to maximise the data transmitted from the two gossamer ring passages, the read-out cycle was set to the fastest useful mode during the respective passage. For the ring passage on 5 November 2002 this meant that dust data were read-out from the instrument memory and written to the Galileo tape recorder in so-called record mode which started at 02:44 UTC, i.e. 18 min before Galileo crossed Io's orbit during approach to Jupiter. The latest data set measured in each amplitude range was read-out at approximately 1 min intervals and written to the onboard tape recorder for later transmission to Earth. Hence, for impact rates up to  $\sim 1 \text{ min}^{-1}$  in each amplitude range, all data sets could be transmitted to Earth. For higher rates, a fraction of these data sets were lost. This mode gave the highest time resolution of the dust measurements at any time during the mission: about 1 min. The completeness of the transmitted data sets varied between 100% in the highest amplitude ranges (AR2–4) in the faint ring extension beyond Thebe's

orbit down to only 4% for the lowest amplitude range (AR1) in the more populated Amalthea ring.

Dust data were obtained in record mode during Galileo's approach to Jupiter until a spacecraft anomaly (safing) on 5 November 2002 at 06:35 UTC prevented the collection of further data. This anomaly occurred at a distance of  $2.33R_J$  from Jupiter, 16 min after closest approach to Amalthea (at  $2.54R_J$ ) and limited the total period of dust measurements obtained from the gossamer rings to about 100 min. Although the instrument continued to measure dust impacts after the spacecraft anomaly, the data were not written to the tape and, hence, most of them were lost. Only the data sets of a few impact events which occurred in the ring region traversed by Galileo after the spacecraft anomaly were obtained from a full memory readout on 18 November 2002. These data, however, have only a low time resolution of about 4.3 h which is on the order of the duration of the entire gossamer ring passage. Only the total number of events (dust plus noise) in each amplitude range can be derived from the accumulators for the ring region traversed after the spacecraft anomaly.

During Galileo's second gossamer ring passage on 21 September 2003, the dust data had to be transmitted to Earth immediately because the spacecraft struck Jupiter and was destroyed less than an hour later. Therefore, the dust instrument memory was read-out in the fastest mode that allowed data to be transmitted in real time (realtime science mode; see Krüger et al. (2001)). Unfortunately, time resolution in this mode was limited to 7 min. The overall completeness of the transmitted data was about 10% in the faint Thebe ring extension and about 5% in the Thebe ring. Due to lower count rates in the higher amplitude ranges, the completeness of transmission was generally better in the higher amplitude ranges. The last data set from the Galileo dust instrument received on Earth was read out from the dust instrument memory at 17:59 UTC when the spacecraft was at a joventric distance of about  $2.5R_J$ . Thus, data from this ring passage provided in-situ dust measurements from the gossamer rings for a total period of about 60 min with no measurements coming from within Amalthea's orbit.


The motion of Galileo through the gossamer rings together with the readout frequency of the dust instrument memory defined the maximum spatial resolution achievable with the ring measurements. During the first ring passage, with 1 min readout frequency in record mode, Galileo moved  $\sim 1800$  km through the ring along its trajectory between two adjacent instrument readouts. This corresponds to a motion in radial distance of about 1100 km (or  $0.015R_J$ ). For the second ring passage the radial resolution was only about 14,000 km or  $0.2R_J$ . The ring and the Galileo trajectory are sketched in Figs. 1 and 2 and the characteristics of both ring passages are summarized in Table 1.

During the entire first ring passage a total of several thousand dust impacts were counted. Approximately 330 of these happened before the spacecraft safing at  $2.33R_J$  inbound to Jupiter. With the optimised noise identification scheme described in Section 2.2 complete data sets of 90 true dust impacts were identified in the Galileo recorded data from the region between  $3.75R_J$  and  $2.33R_J$ . During the second ring passage approximately 260 dust impacts were counted down to  $2.5R_J$  inbound to Jupiter. At this distance dust data transmission ceased before Galileo hit Jupiter. Twenty data sets of dust impacts detected between  $3.75R_J$  and  $2.5R_J$  were transmitted to Earth.

### 2.4. Mass and speed calibration

Grain impact speeds and masses are usually derived from Eq. (1) and an empirical calibration obtained in the laboratory (Grün et al., 1995). Analysis of the dust data measured during Galileo's entire Jupiter mission, however, revealed strong degradation of the instrument electronics which affected the speed and mass calibration. The degradation was most likely caused by the harsh radiation environ-





**Fig. 1.** Projection of Galileo's trajectory through the gossamer ring region on 5 November 2002 (Galileo orbit A34) and 21 September 2003 (orbit J35) onto Jupiter's equatorial plane. The orbits of the small moons Thebe, Amalthea, Adrastea and Metis are indicated by dashed lines. Crosses indicate 1-h time intervals. The different gossamer ring regions are highlighted. Galileo's closest approach to Amalthea occurred on 5 November 2002 at 06:19 UTC (indicated by a filled circle). Thick solid sections of Galileo's trajectory indicate time periods when dust data were obtained.

ion collector lead to prolonged rise time measurements, (iv) degradation of the channeltron required five increases of the channeltron high voltage during the Galileo Jupiter mission, (v) no impact or noise event was registered in the highest ion charge amplitude ranges AR5 and AR6 after July 1999. In particular, (ii) and (iii) affect the mass and speed calibration of the dust instrument. For dust measurements taken after the year 2000, masses and speeds derived from the instrument calibration must be taken with caution because the electronics degradation was severe. Only in cases where impact speeds are known from other arguments, such as exist here in the gossamer rings, can reliable particle masses be derived. This will be discussed in more detail in Section 3.3.

### 3. Results


#### 3.1. Dust impact rates

In Fig. 5 we show examples of the impact rates measured during either gossamer ring passage of Galileo as derived from the accumulators of the dust instrument. We show the rates for the classes and amplitude ranges for which a sufficiently large number of events were counted so that meaningful rate curves could be derived.


The rates measured in all categories (i.e. classes and ion amplitude ranges) increased during approach to Jupiter. From the outer edge of the Thebe ring extension until the time when the dust measurements stopped in the Amalthea ring due to the spacecraft anomaly, the increase was about two orders of magnitude in the lowest channels, AR1, whereas it was only one order of magnitude in the higher channels (AR2–4). This indicates a higher fraction of small particles in the Amalthea ring than in the Thebe ring and the faint Thebe ring extension. In all channels, the highest rates occurred inside Amalthea's orbit when the spacecraft crossed into the more densely populated Amalthea ring. No impacts were measured in the largest categories AR5 and AR6 during both gossamer ring passages.

The instrument accumulators do not contain any information of whether the counted events were due to noise or real dust impacts.

ment in the inner jovian magnetosphere, and a detailed analysis was published by Krüger et al. (2005). Here we recall only the most significant results which are relevant for the gossamer ring measurements: (i) the sensitivity of the instrument for dust impacts and noise dropped with time, (ii) the amplification of the charge amplifiers degraded, leading to reduced measured impact charge values, (iii) drifts in the charge rise times measured at the target and the



**Fig. 2.** Top: Mosaic of Galileo images of Jupiter's gossamer rings taken when the spacecraft was very nearly in the ring plane (from Burns et al. (1999)). The halo and main ring are overexposed (solid white) at the left hand side of the image. To the right are the Amalthea ring (shown in yellow) and the Thebe ring (shown in red). Crosses mark the four extremes of the radial and vertical motions of Amalthea and Thebe as caused by their eccentric and inclined orbits. A very faint extension reaches out beyond Thebe's orbit (shown in blue). Bottom: Galileo's trajectories during the ring passages on 5 November 2002 (solid line) and 21 September 2003 (dashed line). The sections where dust data were collected during both passages are highlighted as thick lines. The approximate locations of the moons' orbits are indicated by vertical dashed lines and Amalthea's position during closest approach on 5 November 2002 is marked by a filled circle (from Hamilton and Krüger (2008)) (for interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article).



**Fig. 3.** Galileo spacecraft configuration (schematic). *Top:* Side view; *Bottom:* Top view. The dust detector (DDS) is mounted directly underneath the magnetometer (MAG) boom (Kivelson et al. (1992)). The sensor field-of-view (FOV) is shown by dashed lines. The locations of the Plasma Instrument (PLS) (PLS; Frank et al., 1992) and the Energetic Particles Detector (EPD) (EPD; Williams et al., 1992), which partially obscure the DDS FOV, are also indicated.


**Table 1**  
Characteristics of Galileo gossamer ring dust measurements.

Date (Galileo orbit number)	5 November 2002 (A34)	21 September 2003 (J35)
Distance range measured	$>2.33R_J$	$\geq 2.5R_J$
Measurement time within $3.75R_J$	100 min	60 min
Time resolution	1 min	7 min
Spatial resolution (radial)	$0.015R_J$	$0.2R_J$
Number of dust impacts counted	$\approx 330$	$\approx 260$
Number of dust data sets transmitted	90	20
Dust impact speed <sup>†</sup>	$18\text{--}20 \text{ km s}^{-1}$	$26\text{--}30 \text{ km s}^{-1}$
Dust detection threshold	$\sim 0.2 \mu\text{m}$	$\sim 0.2 \mu\text{m}$

<sup>†</sup> Dust particles were assumed to orbit Jupiter on circular prograde uninclined orbits.

Since several of the instrument channels were sensitive to noise (cf. Section 2.2) an empirical noise correction factor had to be applied. This factor can only be derived from the data sets transmitted with their full information and it is taken as the ratio between the number of noise events and the total number of events transmitted within a given time interval (dust plus noise; see also Krüger et al. (2001)). Here, the noise rate was calculated as the average over a 1 h interval. The criteria for the identification of individual noise events in the gossamer ring data are given in Section 2.2.

The rate data from the first ring passage show a dip between Thebe's and Amalthea's orbits for both low and high amplitude ranges. The dip is most clearly evident in the lowest amplitude range AR1 where we have the highest number of counted events. The event rate dropped by about a factor of two to five at this location, and the measurements obtained for other particle sizes and during the second ring passage are all consistent with the existence of this dip. One worry, however, is that the noise rate for classes 1 and 2 and the lower amplitude ranges exceeded 80% during some periods of the ring passage, causing the noise removal to lead to large uncertainties in the impact rate. But class 3, our highest quality class, was noise-free and amplitude range AR4 is also expected to be relatively noise-free in class 2. Focusing our attention to these higher quality data, we also see evidence for a dip, which we plot in the right two panels of Fig. 5. Unfortunately, the event rate detected in class 3 for the A34 passage was too low to produce the corresponding plot, and so we plot the lower quality class 2 in-



**Fig. 4.** Dust instrument FOV and obscuration by the magnetometer boom, the PLS and the EPD instruments for an imaginary observer looking outward from the center of the sensor target. *Left:* first ring passage on 5 November 2002 (A34); *right:* second passage on 21 September 2003 (J35). Concentric circles denote the angular distance  $\phi$  (impact angle) from the sensor axis in  $10^\circ$  steps. The spacecraft spin axis is at  $\phi = 60^\circ$  towards the bottom (marked by a cross). The shaded areas show the modelled range scanned by ring particles on circular prograde orbits during each ring passage (Moissl, 2005). The width of the shaded areas is due to the variation of the angle  $\psi$  between the impact velocity and the anti-Earth spin axis during the motion of Galileo through the ring. Note that the sensor side wall is not considered here.

stead. Additional support for this interpretation comes from recent ring imaging de Pater et al., 2008 and increased energetic particle fluxes measured in the dip region with the EPD instrument onboard Galileo (Norbert Krupp, private communication). We therefore conclude that the dip in the impact rate is real, implying a true drop in the dust number density in the Thebe ring. The consequences for grain dynamics and the ring structure will be discussed in Section 4.4.

An additional feature is the extension of the outer gossamer ring far beyond its previously known outer edge at  $3.75R_J$ . Interestingly, the impact rate profile for the smallest particles is relatively flat beyond  $3.75R_J$  whereas inside this distance it increases towards Jupiter. These small submicron particles do not scatter light well and so cannot be seen in optical images; some may be in the process of escaping the gossamer rings as predicted by Hamilton and Burns (1993).

During its first ring passage on 5 November 2002 Galileo had a close flyby of 100-km Amalthea at a closest approach distance of 244 km from the moon's center. Because the Amalthea gossamer ring is believed to be maintained by collisional ejecta from Amalthea itself, an increased dust impact rate is to be expected in the close vicinity of this moon: Galileo detected ejecta dust clouds within the Hill spheres of all four Galilean moons, but outside the Hill spheres there was no noticeable enhancement (Krüger et al., 1999c; Krüger et al., 2003). Taking the recently determined mass of Amalthea (Anderson et al., 2005), its Hill radius is  $r_{\text{Hill}} = 130$  km, only slightly larger than the moon itself. Thus, Galileo did not cross Amalthea's Hill sphere. A spike in the dust flux was not expected, and is not apparent in the  $\sim 40$ -s period when


Galileo was within 500 km of Amalthea. Determining the role of Amalthea as both a source and sink for gossamer ring dust grains requires detailed physical models of (i) the interplanetary impactor population and (ii) ring particle dynamics. This primarily theoretical task is beyond the scope of the current paper.

### 3.2. Grain impact direction

Images of the gossamer rings taken with Galileo and Earth-based telescopes imply that the orbits of the ring particles have very low inclinations with respect to Jupiter's equatorial plane below  $1.5^\circ$ , and that the majority of the grains move on low-eccentric or even circular orbits (de Pater et al., 1999; Ockert-Bell et al., 1999; Burns et al., 1999). In order to calculate the impact direction of the measured ring particles onto the sensor target and the corresponding effective sensor area for these grains, we assumed that the particles orbit Jupiter on circular prograde trajectories with effectively zero inclination.

The only additional parameters necessary are the spacecraft trajectory (state vectors) and spacecraft orientation. The spacecraft trajectory is shown in Figs. 1 and 2, and the spacecraft orientation is constrained by the fact that the antenna pointed within  $3^\circ$  of the Earth direction during both passages of Galileo through the gossamer rings.

With these assumptions, for particles assumed to be on prograde circular orbits, we calculated the dust impact direction and the corresponding sensor area. During the first ring passage, the angle with respect to the spin axis  $\psi$  varied by only  $4^\circ$  in the time interval of interest here when we obtained high-rate recorded data




**Fig. 5.** Dust impact rates (solid curves) measured during both ring passages as derived from the dust instrument accumulators. The ring passage, class and amplitude range are given for each panel. For the first ring passage (A34) data were smoothed with a boxcar average over three data points while no smoothing was applied to data from the second passage (J35). Vertical dotted lines indicate the average locations of Amalthea ('Am') and Thebe ('Th') along their slightly eccentric orbits (cf. Fig. 2), and the edge of the faint ring extension as seen on images ('Ring Edge'). For the first ring passage (A34) a vertical dashed line indicates the actual location of Amalthea during Galileo's closest approach to this moon. Uncertainties due to both noise removal (where necessary—cf. Section 3.1) and statistical fluctuations within a 10–20 min time interval are indicated by vertical error bars. This averaging time interval is indicated by horizontal error bars.

456 from the ring region. In this interval the target area, averaged over  
457 one spacecraft spin revolution (and ignoring obscuration by the  
458 magnetometer boom, PLS and EPD), was 50–55 cm<sup>2</sup>. During the  
459 second ring passage  $\psi$  varied by about 10° and the sensor target  
460 area changed between 200 and 230 cm<sup>2</sup>. This, as well as the differ-  
461 ence in spacecraft radial speed (Table 1), accounts for the factor of  
462  $\approx 5$  increased flux in the second ring passage (J35) as compared to  
463 the first (A34)—see Fig. 5. For both passages the expected rotation  
464 angle for particles orbiting Jupiter on prograde circular trajectories  
465 was  $\Theta \approx 90^\circ$ , and that for retrograde trajectories  $\Theta \approx 270^\circ$ .


466 The range of the rotation angle distribution  $\Delta\Theta$  is determined  
467 by the sensor FOV which is nominally 140°. A smaller FOV was  
468 found for a subset of the 10-nm-sized jovian dust stream particle  
469 impacts (Krüger et al., 1999b); we believe that this reduction is  
470 due to the small sizes and rapid speeds of stream particles. In the  
471 gossamer rings, by contrast, we expect a larger than nominal effective  
472 FOV; recent analysis of Galileo and Ulysses dust data showed  
473 that the sensor FOV for particles much larger than the jovian dust  
474 streams population is almost 180° because the inner sensor side  
475 wall showed a sensitivity for dust impacts comparable to that of  
476 the target itself (Altobelli et al., 2004; Willis et al., 2004; Willis  
477 et al., 2005). We therefore consider an extended FOV for the analysis  
478 of gossamer ring particles. The rotation angles  $\Theta$  of the dust  
479 impacts measured during both ring passages are shown in Fig. 6  
480 and histograms showing the number of impacts per rotation angle  
481 bin are given in Figs. 7 and 8. The rotation angle distribution measured  
482 during the first ring passage (A34 on 5 November 2002)  
483 shows a broad gap at  $\Theta \approx 90^\circ$  having a width  $\Delta\Theta \approx \pm 20^\circ$ . This is  
484 due to shadowing by the magnetometer boom (see Fig. 4). No such  
485 gap in the distribution occurred during the J35 encounter (Fig. 8)  
486 consistent with the geometry of that final ring passage (Fig. 4).

487 As can be seen in Fig. 6, the distribution of the rotation angles  
488 measured during the first gossamer ring passage is much wider  
489 than expected for a sensor target with 140° FOV.

490 The expected width of the rotation angle distribution for particles  
491 on prograde uninclined circular orbits was  $\Delta\Theta \approx 100^\circ$  (cf.  
492 Fig. 4; an analysis of  $\Delta\Theta$  vs.  $\psi$ —the angle between the impact  
493 direction and the spacecraft spin axis—is given by Krüger (2003,  
494 his Fig. 2.7b). Hence, the distribution of measured rotation angles  
495  $\Theta$  should cover the range  $40^\circ \lesssim \Theta \lesssim 140^\circ$ . About half of the im-  
496 pacts, however, were detected with rotation angles  $\Theta \gtrsim 140^\circ$  or




**Fig. 7.** *Top panel:* Distribution of rotation angles  $\Theta$  measured during the first gossamer ring passage (A34; thick black line) and a modelled detector sensitivity function (thin red line). The model curve takes into account the sensitivity of the dust detector target plus side wall, shading by the magnetometer boom and the PLS and EPD instruments (cf. Fig. 4), and a distribution of the orbit inclinations up to 60°: more than 50% of the detected grains are consistent with inclinations below 10°, about 90% of the inclinations are below 30°, and the maximum inclination is 60° Moissl, 2005[from]. *Bottom panel:* Modelled detector sensitivity functions for particle inclinations  $i = 0^\circ, 30^\circ$  and  $60^\circ$ . Arrows indicate the regions where the magnetometer boom (MAG) and the PLS and EPD instruments obscure the detector FOV. The sensitivity curves are in arbitrary units. The model curves in all panels include the FOV of both the target and the sensor side wall and they are integrated over the entire gossamer ring passage (distance range 3.75–2.33R<sub>J</sub>) (for interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article).



**Fig. 6.** Rotation angles  $\Theta$  of dust impacts measured during both gossamer ring passages. *Left panel:* First ring passage on 5 November 2002 (A34). *Right panel:* Second ring passage on 21 September 2003 (J35). Only impacts are shown for which the complete set of measured impact parameters was transmitted to Earth. Solid nearly horizontal lines indicate the expected width of the rotation angle distribution  $\Delta\Theta$  for circular uninclined particle orbits and a sensor target 140° FOV, while dashed lines show the same for target plus sensor side wall (180° FOV). Vertical dotted lines indicate the orbits of Thebe ('Th') and Amalthea ('Am') and the edge of the faint ring extension as seen on images ('Ring Edge'). We ignore the 1.3° inclination of Jupiter's orbital plane with respect to the ecliptic plane and Jupiter's obliquity of about 3° and take the planet's equatorial plane to be coplanar with the ecliptic plane for simplicity. The symbol sizes indicate the different amplitude ranges (AR1–AR4).





**Fig. 8.** Distribution of rotation angles  $\Theta$  measured during the second gossamer ring passage (J35). The vertical lines indicate the expected width from the FOV for the target and the detector side wall for a distribution of particles on uninclined orbits.

497  $\Theta \lesssim 40^\circ$ . If we include the sensor side wall, the expected range  
498 widens to  $\Delta\Theta \simeq 160^\circ$  but is still smaller than the measured range.  
499 A similarly broad distribution was also measured during the second  
500 ring passage on 21 September 2003.

501 The rotation angle distribution shows even more structure than  
502 just the gap at  $\Theta \simeq 90 \pm 20^\circ$ : Fig. 7 (top panel) reveals an asymme-  
503 try in the sense that the distribution with rotation angles  $\Theta \geq 90^\circ$   
504 is broader and shallower than the one with  $\Theta \leq 90^\circ$ . Moissl (2005)  
505 modelled the detector sensitivity and shadowing of the dust sensor  
506 FOV by the magnetometer boom (MAG), the PLS and EPD instru-  
507 ments. The model assumes an inclination distribution consistent  
508 with the measured rotation angles (Fig. 6) and a sensitive area of  
509 target and side wall.

510 A model curve for particles on circular joventric orbits with up  
511 to  $60^\circ$  inclinations is shown as a red solid line in the top panel of  
512 Fig. 7. It gives an overall good agreement with the measured distri-  
513 bution, in particular considering that the spacecraft structures  
514 shading the dust sensor are described by relatively simple approxi-  
515 mations and that the statistics of detected grains is rather low.  
516 Deviations occur at  $\Theta \sim 60 \pm 10^\circ$  and at the edge of the dust sensor  
517 FOV at  $\Theta \gtrsim 170^\circ$ . In both cases the model underestimates the true  
518 number of detections. It has to be noted that particularly large  
519 uncertainties occur at the edge of the FOV where the sensitive area  
520 drops to zero. Also, the modelled curve underestimates the true  
521 width  $\Delta\Theta$  of the rotation angle distribution. It indicates that a lar-  
522 ger fraction of the detected grains may have had orbits with incli-  
523 nations up to about  $60^\circ$  and eccentricities up to 0.2 (Moissl, 2005).  
524 In all, the particle orbits significantly differ from the circular unin-  
525 clined case implied by the ring images.

526 In order to illustrate the significance of the orbital inclinations  
527 on the width  $\Delta\Theta$  of the rotation angle distribution we show in  
528 the bottom panel of Fig. 7 the detector sensitivities averaged over  
529 the entire ring passage (distance range  $3.75$  to  $2.33R_J$ ) for dust par-  
530 ticles with three single inclinations:  $i = 0^\circ$ ,  $30^\circ$  and  $60^\circ$ . As ex-  
531 pected, the rotation angle distribution becomes wider with  
532 increasing inclinations. Note that the sensitivity for dust detections  
533 from certain narrow rotation angle ranges dropped to zero due to  
534 shading by PLS and the magnetometer boom (indicated by arrows)  
535 while EPD obscured the dust beam only during a fraction of the en-  
536 tire ring passage (cf. Fig. 4) so that the sensitivity towards the  
537 direction of EPD is reduced but not to zero.

538 One additional potential reason for the extended rotation angle  
539 distribution may be impacts onto the spacecraft structure close to  
540 the dust sensor. Impacts preferentially onto the magnetometer

boom may have generated impact plasma and secondary grain  
498 fragments which may have hit the dust sensor, resembling true im-  
499 pacts at rotation angles where direct impacts of ring particles onto  
500 the target are impossible. Such events should have revealed their  
501 presence by peculiar impact parameters (charge amplitudes, rise  
502 times, coincidences etc.). An analysis of the data from both ring  
503 passages, however, did not show evidence for such peculiarities  
504 for the majority of grains, making this explanation unlikely (Moissl,  
505 2005). The extended distribution appears, therefore, to be due to  
506 the actual distribution of dust and implies large inclinations  
507 ( $i \lesssim 60^\circ$ ) and non-zero eccentricities ( $e \lesssim 0.2$ ) for many dust par-  
508 ticles. Inclinations and eccentricities of this magnitude are expected  
509 from the model of Hamilton and Krüger (2008, cf. Section 4.4).  
510

### 511 3.3. Grain masses

512 About 90% of the dust impacts measured during both gossamer  
513 ring passages showed abnormally long rise times of the impact  
514 charge signal caused by degradation of the instrument electronics  
515 (Section 2.4). Application of the instrument calibration derived in  
516 the laboratory before launch would lead to unrealistically low im-  
517 pact speeds and, consequently, erroneously large grain masses.  
518 Thus, the rise time measurement cannot be used for calculating  
519 grain impact speeds. In the gossamer rings, impact speeds are  
520 dominated by the spacecraft's speed and, assuming that the parti-  
521 cles move on nearly uninclined circular orbits, the impact speed  
522 onto the detector target on 5 November 2002 was about  $18 \text{ km s}^{-1}$ .  
523 We use this fact as the basis for a procedure to obtain the particle  
524 mass and the number density distributions in this and the follow-  
525 ing section. An overview of the individual processing steps is given  
526 in Fig. 9.

527 We begin by taking  $18 \text{ km s}^{-1}$  instead of the speed derived from  
528 the rise time measurement and calculate the particle mass with Eq.  
529 1, i.e. employing the linear dependence between particle mass  $m$   
530 and impact charge  $Q$ . Similar mass calibration methods were suc-  
531 cessfully applied to earlier measurements of interstellar dust  
532 grains (Landgraf et al., 2000) and to dust impacts measured in  
533 the vicinity of the Galilean moons (Krüger et al., 2000, 2003).  
534

535 An extra complication here is the amplifier degradation that ar-  
536 ose from the accumulated radiation damage to the dust instru-  
537 ment. The damage causes the measured charge amplitude  $Q$  to  
538 be too low by a time-dependent factor that has been calculated  
539 by Krüger et al. (2005). For the time period of interest, we estimate  
540 the additional radiation damage received by the spacecraft and  
541 determine a correction factor of 5 for the ion collector channel  
542 and a factor of 2 for the electron channel, respectively. This means  
543 that measured charges for gossamer ring particles need to be in-  
544 creased by a factor of 5 and 2, respectively, to determine the true  
545 impact charges for these channels. Due to the linear dependence  
546 between impact charge and grain mass (Eq. (1)) this leads to an  
547 average shift in grain mass by a factor of 3.5.  
548

549 In Fig. 10 we show the mass distributions derived for four dif-  
550 ferent regions of the gossamer rings. We include measurements  
551 from: (i) the region between Io's orbit and the outer edge of the  
552 Thebe Extension ( $6$ – $3.75R_J$ ), (ii) the Thebe Extension (between  
553  $3.75R_J$  and Thebe's orbit), (iii) the Thebe ring (between Thebe's  
554 and Amalthea's orbit), and (iv) the Amalthea ring (inside Amal-  
555 thea's orbit). Dust in the outermost of these regions is poorly sam-  
556 pled by the spacecraft and invisible from the ground. Better  
557 statistics exist for dust amongst the Galilean satellites (Grün  
558 et al., 1998; Thiessehusen et al., 2000; Krivov et al., 2002a,b; Zee-  
559 handelaar and Hamilton, 2007).  
560

561 To illustrate the significance of the corrections for instrument  
562 aging and for incomplete data transmission, we show both uncor-  
563 rected and corrected histograms. The aging correction shifts the  
564 entire distribution by a factor of 3.5 to higher masses. Coinciden-






Fig. 9. Flow chart illustrating the individual processing steps required to derive mass and number density distributions for the gossamer rings.

tally, this corresponds to the width of half an amplitude range interval on a logarithmic scale so that the aging correction shifts the mass distribution by one histogram bin. Furthermore, to correct for incomplete transmission, we calculated a correction factor from the ratio between the number of counted impacts and the number of data sets transmitted in a given time interval. We took into account that the leftmost two bins correspond to AR1, the next two bins to AR2 and so on. Note that the transmission correction is most significant in the leftmost two bins (AR1) and nearly negligible in the other bins.

According to Fig. 10 the largest detected particles have masses  $m \approx 5 \times 10^{-13}$  kg. Assuming spherical particles with density  $\rho = 1000 \text{ kg m}^{-3}$  (representative of water ice), the corresponding grain radius is  $s \approx 5 \mu\text{m}$ . For grain densities of 500 and 2000  $\text{kg m}^{-3}$  the grain radius is 6 and 4  $\mu\text{m}$ , respectively. Similarly, the smallest mass just exceeding the detection threshold,  $m \approx 5 \times 10^{-17}$  kg, corresponds to  $s \approx 0.2 \mu\text{m}$ . Thus,  $0.2 \mu\text{m} \leq s \leq 5 \mu\text{m}$  is a plausible size range from the calibration of the impact charges after correction for electronics aging. This shows that the size distribution extends to particles one order of magnitude smaller than derived from ring images. On the other hand, the largest sizes agree rather well with particle sizes deduced from imaging of the gossamer ring (Showalter et al., 1985; Showalter et al., 2008; de Pater et al., 2008) and Jupiter's main ring (Throop et al., 2004; Brooks et al., 2004). The only other information on ring particle sizes comes from three impacts detected at ring plane crossing by the Pioneer 10 and Pioneer 11 spacecraft Humes, 1976. The Pioneer 10 detector was sensitive to particles larger than about 6  $\mu\text{m}$  while the Pioneer 11 detector was sensitive to particles roughly twice as large; these early measurements first showed that there was 10 micron dust in Jupiter's equatorial plane.


Only 20 data sets of impact events were transmitted from the second ring passage (J35) and this low number does not allow us to derive statistically meaningful mass distributions for the individual ring regions. In addition, the mass calibration of these data is even more uncertain because of the rapid degradation of the dust instrument electronics due to accelerated radiation damage very close to Jupiter (Krüger et al., 2005, their Fig. 2).

It is evident that the mass distribution is very similar in the faint Thebe ring extension and in the Thebe ring, while it is much steeper in the Amalthea ring. One has to keep in mind, however, that this steeper slope is dominated by the leftmost two bins of the distribution for masses  $\sim 5 \times 10^{-16}$ – $5 \times 10^{-17}$  kg which required the largest corrections for noise removal and incomplete transmission. Although these bins required the largest corrections we are convinced that the strong excess in small grains is real.

The slopes of the incremental mass distributions (Colwell, 1993) given by  $\Delta \log N(m) / \Delta \log m \propto m^\gamma$  (with  $N(m)$  being the number of particles per logarithmic mass interval) for the individual ring regions are listed in Table 2. While the slopes of the Thebe ring and Thebe extension are well reproduced by power laws the slope for the Amalthea ring is poorly described by a power law.

Note that in all histograms the leftmost bin is lower than the next one at higher masses. This is a well-known effect (Krüger et al., 2006, their Fig. 6) and is most likely due to the fact that the sensitivity threshold of the dust instrument may not be sharp. We therefore did not include the leftmost bin in the fitting of power law slopes to the mass distributions.

Interestingly, the slopes tend to steepen significantly when going from the outer to the inner ring regions (although ignoring the two leftmost bins in the distribution for the Amalthea ring would make this distribution similar to the one for the Thebe ring). This is due to the weakening of electromagnetic forces in the vicin-



**Fig. 10.** Incremental mass distributions per logarithmic mass interval for different regions of the gossamer rings obtained from the first ring passage on 5 November 2002 (A34; 87 particles with velocity error factor VEF <6; Grün et al., 1995). An impact speed of  $18 \text{ km s}^{-1}$  was assumed to calculate grain masses from the measured charge amplitudes (Eq. (1)). The detection threshold for the assumed impact speed is indicated by vertical dotted lines (without instrument aging), and by vertical dashed lines (with aging correction). Dotted histograms show the distribution with neither corrections for instrument aging nor incomplete data transmission. The solid histograms show the distribution corrected for both incomplete data transmission (which increases the number in the dotted bins) and instrument aging (which shifts the bins to the right). The thick solid lines are linear fits to these corrected histograms. The linear fits seem fine for all but the Amalthea ring. The slopes for the mass distributions are given in Table 2.

ity of synchronous orbit ( $2.25R_J$ )—small particles that can be expelled from the Thebe ring cannot be ejected from the Amalthea ring (Hamilton and Burns, 1993; Hamilton and Krüger, 2008) leading to enhanced number densities there.

The power law slopes obtained for the individual ring regions agree very well with the slopes measured in-situ in impact-generated dust clouds at the Galilean moons (Krüger et al., 2003), while they are much flatter than slopes derived for Saturn’s E ring (Kempf et al., 2008). This indicates that the majority of the detected grains may be collisional ejecta from hypervelocity impacts onto the surfaces of parent bodies embedded in the gossamer rings (mostly Amalthea and Thebe).

### 3.4. Dust number density

Each of the impact charge amplitude ranges of the dust instrument corresponds to a factor of 10 in impact charge and, hence, a factor of 10 in mass (for constant impact speed; cf. Eq. (1)). Therefore, a number density distribution derived from the accumulators directly reflects the grain mass distribution. We use this approach to construct relative grain size distributions in the individual gossamer rings without using the dust instrument calibration from the laboratory. The individual data processing steps are again summarised in Fig. 9.


The dust number density  $n$  is proportional to the impact rate  $\Delta N/\Delta t$  recorded by the dust instrument, and the relation between both quantities is given by:

$$n = \frac{\Delta N}{\Delta t} \cdot \frac{1}{v \cdot A_S(\psi)} \quad (2)$$

$A_S(\psi)$  is the sensor area as a function of the angle  $\psi$  with respect to the spacecraft spin axis, and  $v$  is the grain impact speed. To obtain impact rates, we separated different ring regions into distance bins and divided the number of particles  $\Delta N$  counted in a given distance bin by the time  $\Delta t$  Galileo spent in this bin.

In Fig. 11 we show the number densities derived from the accumulators of the four amplitude ranges for the individual gossamer ring regions. Number densities measured during both gossamer ring passages agree to within about 50%, except in the region between Io’s orbit and the outer ring edge. Here the measurements disagree by a factor of 3 (Fig. 11). Despite the low number of dust detections in this ring region and the uncertainty due to the noise removal, we believe that this difference in the number density is likely real, pointing to azimuthal variations in the dust ring density itself.

Hamilton and Krüger (2008) have proposed that a shadow resonance governs the behavior of the gossamer rings and their Fig. 3 shows that the diffuse outer Thebe ring should be asymmetric and offset away from the Sun. Such a structure would yield a larger impact flux to a spacecraft approaching from the anti-Sun hemisphere (A34, the first passage) than from the sunward hemisphere (J35, the second passage)—see Fig. 1. This is in qualitative agreement with the difference in the outermost ring regions observed here. Moreover, the Hamilton and Krüger model also pre-



**Fig. 11.** Incremental dust number density distributions per logarithmic mass interval deduced from the accumulators (classes 1–3 taken together after noise removal). Solid lines show the data for the first gossamer ring passage (A34), dotted lines show those for the second passage on 21 September 2003 (J35). For the Thebe ring, the A34 and J35 data agree quite well, while for the Amalthea ring no J35 data were obtained. Data for the distant ring in panel 1 disagree; possible reasons are discussed in the text. The assumed grain impact speeds are  $18\text{--}20\text{ km s}^{-1}$  and  $26\text{--}30\text{ km s}^{-1}$ , respectively. The slopes for these number density distributions are given in Table 2.

**Table 2**

Slopes  $\gamma$  of the mass distributions derived in this work for the different ring regions (1). The Galileo orbits from which these data are derived are indicated. (2) lists the slope of the mass distribution as derived from the instrument calibration (Fig. 10), and (3) and (4) the ones obtained from the measured number densities (Fig. 11), respectively. In column 4 the slope for the region between the outer ring limit and Io's orbit is put in parentheses because it is derived from a very low number of detections. We have put the numbers in column (2) in bold face to emphasize that they are the most reliable.

Population (1)	From calibration	From number density	
	A34 (2)	A34 (3)	J35 (4)
Amalthea ring	<b><math>-0.76 \pm 0.51</math></b>	$-0.42 \pm 0.39$	–
Thebe ring	<b><math>-0.24 \pm 0.13</math></b>	$-0.17 \pm 0.18$	$-0.23 \pm 0.42$
Thebe ring extension	<b><math>-0.31 \pm 0.16</math></b>	$-0.22 \pm 0.22$	$-0.20 \pm 0.28$
Io to ring limit	<b><math>-0.09 \pm 0.18</math></b>	$-0.01 \pm 0.09$	$(-0.30 \pm 0.00)$

dicts that larger particles should not spread very far outward from their Thebe and Amalthea sources in agreement with the lack of AR4 grains in Fig. 11 beyond the outer visible edge of the Thebe ring.

Total number densities obtained by adding the values for each histogram bin in each panel are given in Table 3. These values take into account the sensor target only. If we assume that the sensitivity of the side wall is the same as that of the target, the number densities derived from the first ring passage are lower by about 50% while those for the second passage are reduced by only about 10%. This leads to somewhat better agreement between the two

passages. For the mass densities given in Table 3 we have assumed spherical grains with density  $1000\text{ kg m}^{-3}$ .

In Table 3 we also give number densities for dust populations detected by Galileo beyond the orbit of Io. Number densities derived for the various ring regions smoothly drop with increasing jovicentric distance, showing that Jupiter's faint ring system fills the entire space from the gossamer rings close to Jupiter out to the region of the Galilean moons and beyond.

## 4. Discussion

### 4.1. Comparison of in-situ data and remote imaging

From optical imaging, ring particle size distributions can be estimated by making assumptions about grain optical properties including the real and imaginary components of the index of refraction and roughness parameters. Similarly, deriving size distributions from the Galileo dust impact data requires assumptions about instrument aging and impact velocities. When both optical and in-situ data are available, a new hybrid method for determining sizes is possible.

The new method has the advantage of depending only on well-measured quantities: the ring normal optical depth,  $\tau$ , the ring's vertical extension,  $H$ , both derived from imaging, and the number density,  $n$ , measured in-situ. In particular, this calculation is independent of the mass calibration of the dust instrument. Relevant ring properties are given in Table 4. The optical depth has the big-



**Table 3**  
Physical parameters of dust populations (1) detected in-situ at Jupiter. (2) lists the radial distance range where the particles were detected, (3) gives typical particle radii assuming spherical particles, (4) and (5) give the derived particle number densities and mass densities in space, respectively, (6) lists the dust mass contained in small particles (0.2–5  $\mu\text{m}$ ), and (7) gives references.

Population (1)	Jovicentric distance ( $R_j$ ) (2)	Particle radii ( $\mu\text{m}$ ) (3)	Number density ( $\text{km}^{-3}$ ) (4)	Mass density ( $\text{kg m}^{-3}$ ) (5)	Dust mass in small grains (kg) (6)	Reference (7)
Amalthea ring	2.33–2.54	0.2–5	$\sim 2 \times 10^6$	$\sim 4 \times 10^{-18}$	$\sim 10^6$	This work
Thebe ring	2.54–3.1	0.2–5	$\sim 3 \times 10^5$	$\sim 10^{-18}$	$\sim 2 \times 10^6$	This work
Thebe ring extension	3.1–3.75	0.2–5	$\sim 10^5$	$\sim 4 \times 10^{-19}$	$\sim 10^6$	This work
Io to ring limit	3.75–6	0.2–2	$\sim 5 \times 10^3$	$\sim 5 \times 10^{-21}$	$\approx 5 \times 10^4$	This work
Galilean ring	10–30	0.6–3	$10^2$ – $10^3$	$10^{-21}$ – $10^{-20}$		Krivov et al. (2002a)
Captured particles	10–20	0.5–1.5	$\sim 10^2$	$\sim 10^{-21}$		Thiessenhusen et al. (2000)
Distant ring	$\geq 50$	1–2	$\sim 10^1$	$\sim 10^{-22}$		Krivov et al. (2002b)

**Table 4**  
Properties of the gossamer rings as obtained from imaging observations (Showalter et al., 1985; Ockert-Bell et al., 1999; de Pater et al., 1999).

	Amalthea ring	Thebe ring	Uncertainty
Normal optical depth $\tau$	$10^{-7}$	$3 \times 10^{-8}$	Factor of 5
Ring half-thickness $H$	1300 km	4400 km	$\pm 100$ km

gest error bar whereas the ring's vertical extension is rather well known. Furthermore, imaging shows that the rings are most tenuous near-Jupiter's equatorial plane and densest near their vertical limits (Ockert-Bell et al., 1999; de Pater et al., 1999). A recent ground-based determination of the optical depth by de Pater et al. (2008) is a factor of 5 above the Showalter et al. (2008) values used here. The latter values are more relevant for our purposes as the particle populations that they probe are closer to those sampled by the dust detector.

The typical ring particle radius can be expressed as

$$s = \sqrt{\frac{\tau}{2\pi H n_{\text{opt}}}} \quad (3)$$

Here,  $n_{\text{opt}}$  is the number density measured in-situ of grains dominating the optical cross-section. But what should we use for  $n_{\text{opt}}$ ? Summing over all amplitude ranges yields the number densities given in Table 3 and an effective grain radius  $s \approx 2 \mu\text{m}$ . In this simple analysis all measured particle sizes contribute to the optical cross-section.

For a more realistic calculation we have to take into account that imaging is most sensitive to those particles which have the largest cross-section for reflecting light. Using the fact that amplitude ranges AR1–4 correspond to a factor of 1000 in mass (100 in area), Fig. 12 shows the relative contribution of the four amplitude ranges to the optical cross-section. In all ring regions the biggest contribution to the optical depth comes from the biggest grains (AR4), even though the smallest ones (AR1) dominate the number density. Thus, a better choice for  $n_{\text{opt}}$  is to use AR4 only.

Now taking the number densities from Fig. 11 for AR4 only, the derived grain radii are  $s \approx 5 \mu\text{m}$  for the Thebe ring and  $\approx 10 \mu\text{m}$  for the Amalthea ring, respectively. In order to estimate the uncertainty in these grain sizes, one has to take into account the uncertainties in the imaging observations and in-situ measurements alike. The uncertainty in the imaging is dominated by that of the optical depth  $\tau$  which is about a factor of 5 (Table 4; the uncertainty of the ring half-thickness  $H$  is below 10%). For the in-situ measurements it is the uncertainty in the highest channels, AR4, which counts here. Given that AR4 needs at most a small noise correction (Section 3.1), we think that the uncertainty in this correction is much less than a factor of two. Adding another factor of two for the statistical uncertainty due to the low number of dust impacts, the overall uncertainty in the number density is about a


factor of 3. This leads to an uncertainty in the grain radii which is about a factor of 4.

These grain sizes derived from Eq. (3) and the number densities in AR4 are consistent with the optical measurements Showalter et al., 2008; de Pater et al., 2008, and they agree within about a factor of 2 with the biggest sizes obtained from the calibrated in-situ data. Given the overall uncertainties of the dust instrument calibration and the calculation of the optical depths, the agreement between all of these methods is quite satisfactory.

An interesting quantity that we can derive from our analysis is the relative contribution of grains on inclined orbits to the number density and, hence, to the optical depth. Figs. 6 and 7 indicate that about 20% of the measured grains are incompatible with uninclined orbits, requiring an inclination of  $20^\circ$  or greater. This implies that in the ring plane these grains contribute about 20% to the total number density of dust larger than approximately  $0.2 \mu\text{m}$ . Their contribution to the optical depth, however, is somewhat lower because most of these grains are sub-micron in size. Fig. 12 shows that the contribution of the sub-micron grains (AR1 and AR2) to the total cross-section is typically about 5%, rising in the Amalthea ring to perhaps 20%. This implies that the grains on inclined orbits contribute on the order of 1–4% to the total optical depth, the larger value being applicable to the Amalthea ring. This small percentage is well below the limits of detectability with today's imaging techniques, especially considering the fact that these grains would be spread over a range approximately 20 times greater than the vertical extent of the Thebe ring.

#### 4.2. Grain size distributions

In Sections 3.3 and 3.4 we determined the grain mass distributions in two different ways. Both analyses produced the steepest distributions in the Amalthea ring while further away from Jupiter the distributions are much flatter. However, the slopes derived from the number density distributions (Section 3.4) are somewhat flatter than those obtained from the mass distributions (Section 3.3, see also Table 2). These flatter slopes are probably due to an unsharp detection threshold of the dust instrument (Krüger et al., 2006), leading to an unrealistically depleted leftmost mass bin for the smallest particles (Fig. 10). In order to get an estimate of the influence of this effect on the slopes derived from the number densities, we recalculated the mass distributions by including all bins in the fit: the mass distributions became flatter, except for the Amalthea ring (see below), and they agreed very well with the slopes derived from the number densities. This supports our contention that the leftmost mass bin is incomplete and should be ignored as we do in our derivation of column 2 of Table 2. We therefore conclude that the slopes of the mass distributions obtained from the instrument calibration are a better measure of the true distributions in the ring than those derived from the number densities.



**Fig. 12.** Normalised relative cross-section of dust particles deduced from the accumulators (class 1–3 taken together). Solid lines show the data for the first gossamer ring passage (A34) and dotted lines show those for the second passage on 21 September 2003 (J35). These panels show that the large AR4 grains dominate the optical depth in all ring components. Note that since the sum of all bins in each histogram is normalised to one, the heights of the solid and dotted histograms cannot be directly compared.

In the Amalthea ring the fit with all bins gives a slope of  $-0.63 \pm 0.43$  which is somewhat steeper than the slope obtained from the number density ( $-0.42 \pm 0.39$ ). This may indicate that the correction for incomplete transmission for the Amalthea ring (which mostly affects the two left-most bins in the mass distribution) is too strong. More likely, this mismatch simply means that the Amalthea distributions are not well fit by simple power law distributions as can be clearly seen in the figures.

Showalter et al. (2008) derived a size distribution for the Amalthea ring which is brightest in imaging. They get a power law slope of  $-2$  to  $-2.5$  in the size range  $4\text{--}30\ \mu\text{m}$ . Therefore, the in-situ measurements and the imaging results complement each other with only little overlap in the sensitive size range. Furthermore, a size distribution for the main jovian ring was recently determined from Galileo observations by Brooks et al. (2004). They find a power law slope of  $-2.0 \pm 0.3$  for particles below  $\sim 15\ \mu\text{m}$  and a transition to a power law with slope  $-5.0 \pm 1.5$  at larger sizes.


In Fig. 13 we compare these distributions with our in-situ measurements. Note that the size distribution for the Amalthea ring derived from our in-situ measurements for the small grains agrees very well with the one obtained from images for large grains. Beyond Amalthea's orbit the size distribution for submicron grains becomes flatter while little is known about the abundance of grains bigger than  $5\ \mu\text{m}$  in these regions.

Fig. 13 is the most complete compilation of the grain size distributions in the jovian ring system presently available. It is obvious that even though the small submicron particles are the most abundant in the rings (top panel), the largest contribution to the total ring mass comes from the bigger grains above  $10\ \mu\text{m}$  (bottom panel; see also Section 4.3).

#### 4.3. Total ring mass

From the number density measured in-situ in the rings (Fig. 11) and the known ring volume, we calculate the entire ring dust mass contained in the small particles ( $0.2\text{--}5\ \mu\text{m}$ ). Taking the dimensions of the Amalthea and Thebe rings given in Table 4 and noting that the average density near the midplane is half that of the vertical extremes, the total mass in each of these two gossamer ring components is a few  $10^6$  kg. Note that here we have assumed a smooth dust distribution inward of the source moons. If we take into account that both the Amalthea (de Pater et al., 2008) and Thebe rings are confined to the regions just interior to their bounding satellites, the derived dust masses become somewhat lower (by a factor of 2–3). For the Thebe ring extension we find a similar value of about  $10^6$  kg of dust, assuming that this ring has the same vertical extension as the Thebe ring itself. The ring masses for the Thebe ring and Thebe ring extension derived from Galileo's two independent ring passages agree to within 15%. For the ring region between the outer edge of the Thebe ring extension and Io's orbit we assumed the same vertical extension as for the Thebe ring extension. Note, however, that there is no optical data available for this region and dynamical simulations show that the ring is likely further extended. Therefore, the derived ring mass of  $\approx 5 \times 10^4$  kg is a lower limit. Furthermore, the two ring passages give results that differ by a factor of three as discussed in Section 3.4. This is probably due to the very asymmetric shape of the outermost ring Hamilton and Krüger, 2008. We collect these numbers in Table 3.

The bottom panel of Fig. 13 shows that the small grains measured in-situ represent only a minor fraction of the total ring mass



**Fig. 13.** Relative grain size distributions per logarithmic size interval for the gossamer rings derived in this work (four solid lines) compared with the Galileo imaging results of Showalter et al. (2008, dashed lines). The Cassini imaging results from the main jovian ring fit by Brooks et al. (2004, dotted lines) are overplotted for reference. The vertical axis is in arbitrary units and the curves are shifted so that they all fit together at 3  $\mu\text{m}$ . In each panel, the solid lines refer to—from top to bottom—the Amalthea ring ('Am'), the Thebe ring ('Th'), the Thebe extension ('Th Ext') and the region between the outer ring edge and Io's orbit ('Io to Ring'), respectively. Finally, we show the Galileo dust data for Amalthea as a histogram since it departs so dramatically from a power law. *Top panel:* relative number density of particles in the ring; *Middle panel:* relative cross-sectional area of the ring; *Bottom panel:* relative ring mass density.

contained in the dust: Assuming that the size distribution for optically visible grains in the size range 4–30  $\mu\text{m}$  measured by Showalter et al. (2008) is valid for all gossamer rings, the total ring mass is increased by a factor of  $\sim 30$  over the values for small particles that we list in Table 3. Similarly, if we take the bimodal size distribution derived for the main jovian ring by Brooks et al. (2004) in the size range 0.1–100  $\mu\text{m}$ , the gossamer ring mass increases by a factor of  $\sim 25$ . There is evidence for even more mass at still greater sizes (Showalter et al., 2008).

4.4. Grain dynamics

The interesting properties of the gossamer rings can be most easily explained with the shadow resonance model of Hamilton and Krüger (2008). We briefly summarize their model here, making explicit connections to the observations that we have discussed above. The shadow resonance is an electromagnetic effect that occurs when a dust grain enters Jupiter's shadow, photoelectric charging by solar radiation switches off, and the grain's electric potential decreases. This leads to an oscillating particle charge due to the switch on and off of photoelectric charging on the day and night side of the planet (shadow resonance). It changes the electro-

magnetic force acting on the particle and results in coupled oscillations of the orbital eccentricity and semimajor axis. The oscillations cause the rings to extend significantly outward, but only slightly inward, of their source moons while preserving their vertical thicknesses. This is exactly what is observed for the Thebe ring extension. Furthermore, it leads to longitudinally asymmetric gossamer rings, offset from the Sun for positive grain charges which may be the cause of the number density differences measured between 6 and 3.75 $R_J$  for the two ring passages (Fig. 11). Furthermore, in the absence of a dissipative drag force, the model implies a lack of material inside a certain distance from Jupiter. If most ring material is reabsorbed by the satellites before drag forces can draw it inward, this would create the gap interior to Thebe that is visible in the rate plots in Fig. 5. de Pater et al. (2008) and Showalter et al. (2008) also see evidence for a dropoff of number density interior to Thebe's orbit.

An additional feature of the Galileo gossamer ring data is the likely detection of particles on high inclination orbits. The possibility that spurious events, such as impacts into the detector wall or the magnetometer boom, masquerade as particles with high inclinations can be most likely ruled out. Searching for a physical explanation, the findings are consistent with grains being driven to large inclinations by the shadow resonance as well (Hamilton and Krüger, 2008). The grains would form a halo of material faint enough to be invisible to imaging, but populated enough to be detected by direct impacts onto the Galileo sensor. Showalter et al. (2008) also see indications for a broadening of the inclinations in the Thebe ring, although only to a few degrees above and below the ring plane. Our size distribution extends to an order of magnitude smaller grains than the smallest grains detected by the images and, thus, the expectation that smaller grains should be more sensitive to the shadow resonance and thus on higher inclination orbits would be consistent with our Galileo in-situ data. One would also expect the smaller grains to show a wider distribution in rotation angles than the bigger ones which they in fact do; the impacts measured in AR4 during the A34 passage can mostly be explained with uninclined circular orbits while the smaller particles of AR1 and AR3 need orbit inclinations up to 20°. The more sparse J35 data do not show this same trend, although this may be at least partially due to poor statistics.

Electromagnetic forces in general and the shadow resonance in particular seem to be crucial for determining the structure and dust transport in Jupiter's tenuous gossamer rings. Because dust from a single source is dispersed just slightly inside but widely outside the source, a similar mechanism may also be responsible for the wide outward extension of Saturn's E ring recently detected with the Cassini dust instrument out to at least 18 $R_S$  (Srama et al., 2006, Saturn radius  $R_S = 60,280$  km). Furthermore a large vertical extension recently seen on Cassini images (Ingersoll et al., 2007) is likely due to similar electromagnetic effects.

5. Conclusions

The Galileo in-situ dust detector made the first successful measurements of submicron and micron-sized dust impacts in Jupiter's gossamer rings during two ring passages of the spacecraft in 2002 and 2003. Dust impacts were measured in all three regions of the gossamer rings which had been previously identified on optical images. The region between Io's orbit and the outer limit of the faint Thebe extension, where the ring is invisible to imaging, was also explored. The data from the two ring passages allow for the first actual comparison of in-situ dust measurements with the properties inferred from inverting optical images.

The measured impact rate profile shows a drop immediately interior to Thebe's orbit and the grain impact directions extend over a significantly wider range than expected for grains moving about



Jupiter on uninclined circular orbits. In fact, inclinations up to 20° nicely explain the measured impact directions for most grains. We investigated the idea that spurious events, such as impacts onto the magnetometer boom, masquerade as particles with high inclinations, and are convinced that such explanations can be ruled out.

The wide range in impact directions can be explained by a shadow resonance caused by varying particle charge on the day and night side of Jupiter, driving particles onto high inclination orbits. They form a halo of material faint enough to be invisible to imaging, but populated enough to be detectable with the Galileo sensor. The faint gossamer ring extension previously imaged to about 3.75R<sub>J</sub> was detected out to at least 5R<sub>J</sub>, indicating that ejecta from Thebe spread much further and particle orbits get higher eccentricities than previously known. Both the gap in the ring and the faint ring extension indicate that the grain dynamics is strongly influenced by electromagnetic forces.

The measured grain sizes range from about 0.2 to 5 μm, their abundance increasing towards smaller particles. Our measurements extend the known size distribution for the gossamer rings by a factor of ten towards smaller particles than previously derived from imaging. Within the measurement uncertainties, particles contributing most to the optical cross-section are about 5 μm in radius, in agreement with imaging results. The grain size distribution is consistent with the majority of grains being generated by hypervelocity impacts onto the surfaces of the moons orbiting Jupiter in the gossamer ring region. While the small particles detected in-situ are the most abundant by number, at least an order of magnitude more mass is contained in particles larger than 5 μm which—because of their large surface areas—also dominate ring images. The size distributions of grains measured in the gossamer rings gradually flatten with increasing distance from Jupiter due to the more efficient electromagnetically-induced escape of more distant grains (Hamilton and Burns, 1993; Hamilton and Krüger, 2008).

The Galileo in-situ measurements obtained throughout the jovian magnetosphere show that the dust densities in Jupiter's faint ring system more or less continuously drop from the region of the gossamer rings close to Jupiter out to the Galilean moons and beyond. While the inner ring regions (1–3.5R<sub>J</sub>) can be clearly seen with imaging techniques, only in-situ spacecraft can presently detect the much fainter dust that permeates near jovian space.

## Acknowledgments

The authors wish to thank the Galileo project at NASA/JPL for effective and successful mission operations. This research was supported by the German Bundesministerium für Bildung und Forschung through Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR, Grant 50 QJ 9503 3). Support by MPI für Kernphysik is also gratefully acknowledged. D.P.H. would like to acknowledge NASA Grant No. NNG06GGF99G for support of this work and the MPI für Sonnensystemforschung for a short-term visitor's grant.

## References

- Altobelli, N., Moissl, R., Krüger, H., Landgraf, M., Grün, E., 2004. Influence of wall impacts on the Ulysses dust detector in modelling the interstellar dust flux. *Planet. Space Sci.* 52, 1287–1295.
- Anderson, J.D., Johnson, T.V., Schubert, G., Asmar, S., Jacobson, R.A., Johnston, D., Lau, E.L., Lewis, G., Moore, W.B., Taylor, A., Thomas, P.C., Weinwurm, G., 2005. Amalthea's density is less than that of water. *Science* 308, 1291–1293.
- Baguhl, M., 1993. Identifikation von Staubeinschlägen in den Daten der Mikrometeoriten-Detektoren an Bord der Raumsonden Ulysses und Galileo. PhD thesis, Ruprecht-Karls-Universität, Heidelberg.
- Baguhl, M., Grün, E., Linkert, G., Linkert, D., Siddique, N., 1993. Identification of 'small' dust impacts in the Ulysses dust detector data. *Planet. Space Sci.* 41, 1085–1098.
- Brooks, S.M., Esposito, L.W., Showalter, M.R., Throop, H.B., 2004. The size distribution of Jupiter's main ring from Galileo imaging and spectroscopy. *Icarus* 170, 35–57.

- Burns, J.A., Hamilton, D.P., Showalter, M.R., 2001. Dusty rings and circumplanetary dust: Observations and simple physics. In: Grün, E., Gustafson, B.A.S., Dermott, S.F., Fechtig, H. (Eds.), *Interplanetary Dust*. Springer Verlag, Berlin, Heidelberg, New York, pp. 641–725.
- Burns, J.A., Showalter, M.R., Hamilton, D.P., Nicholson, P.D., de Pater, I., Ockert-Bell, M.E., Thomas, P.C., 1999. The formation of Jupiter's faint rings. *Science* 284, 1146–1150.
- Colwell, J.E., 1993. Power-law confusion: You say incremental, I say differential. *Lunar Planet. Inst. Conf.* 24, 325–326 (abstract).
- de Pater, I., Showalter, M.R., Burns, J.A., Nicholson, P.D., Liu, M.C., Hamilton, D.P., Graham, J.R., 1999. Keck infrared observations of Jupiter's ring system near Earth's 1997 ring plane crossing. *Icarus* 138, 214–223.
- de Pater, I., Showalter, M.R., Macintosh, B., 2008. Keck observations of the 2002–2003 jovian ring plane crossing. *Icarus* 195, 348–360.
- Frank, L.A., Ackerson, K.L., Lee, J.A., English, M.R., Pickett, G.L., 1992. The plasma instrumentation for the Galileo mission. *Space Sci. Rev.* 60, 283–307.
- Göller, J.R., Grün, E., 1989. Calibration of the Galileo/Ulysses dust detectors with different projectile materials and at varying impact angles. *Planet. Space Sci.* 37, 1197–1206.
- Grün, E., Baguhl, M., Hamilton, D.P., Kissel, J., Linkert, D., Linkert, G., Riemann, R., 1995. Reduction of Galileo and Ulysses dust data. *Planet. Space Sci.* 43, 941–951.
- Grün, E., Fechtig, H., Hanner, M.S., Kissel, J., Lindblad, B.A., Linkert, D., Maas, D., Morfill, G.E., Zook, H.A., 1992. The Galileo dust detector. *Space Sci. Rev.* 60, 317–340.
- Grün, E., Hamilton, D.P., Riemann, R., Dermott, S.F., Fechtig, H., Gustafson, B.A., Hanner, M.S., Heck, A., Horányi, M., Kissel, J., Kivelson, M., Krüger, H., Lindblad, B.A., Linkert, D., Linkert, G., Mann, I., McDonnell, J.A.M., Morfill, G.E., Polansky, C., Schwehm, G.H., Srama, R., Zook, H.A., 1996. Dust measurements during Galileo's approach to Jupiter and Io encounter. *Science* 274, 399–401.
- Grün, E., Krüger, H., Graps, A., Hamilton, D.P., Heck, A., Linkert, G., Zook, H., Dermott, S.F., Fechtig, H., Gustafson, B., Hanner, M., Horányi, M., Kissel, J., Lindblad, B., Linkert, G., Mann, I., McDonnell, J.A.M., Morfill, G.E., Polansky, C., Schwehm, G.H., Srama, R., 1998. Galileo observes electromagnetically coupled dust in the Jovian magnetosphere. *J. Geophys. Res.* 103, 20011–20022.
- Hamilton, D.P., Burns, J.A., 1993. Ejection of dust from Jupiter's gossamer ring. *Nature* 364, 695–699.
- Hamilton, D.P., Krüger, H., 2008. Jupiter's shadow sculpts its gossamer rings. *Nature* 453, 72–75.
- Humes, D.H., 1976. The Jovian meteoroid environment. In: Gehrels, T. (Ed.), *Jupiter*. Univ. of Arizona Press, Tucson, pp. 1052–1067.
- Ingersoll, A.P., Dyudina, U.A., Ewald, S.P., and Hedman, M., 2007. Possible Dust Extended From Saturnian Ring Plane or Zodiacal Light as Seen by Cassini ISS. AGU Fall Meeting, B1294+ (abstract).
- Kempf, S., Beckmann, U., Moragas-Klostermeyer, G., Postberg, F., Srama, R., Economou, T., Schmidt, J., Spahn, F., Grün, E., 2008. The E ring in the vicinity of Enceladus. I. Spatial distribution and properties of the ring particles. *Icarus* 193, 420–437.
- Kivelson, M.G., Khurana, K.K., Means, J.D., Russell, C.T., Snare, R.C., 1992. The Galileo magnetic field investigation. *Space Sci. Rev.* 60, 357–383.
- Krivov, A.V., Krüger, H., Grün, E., Thiessenhusen, K.-U., Hamilton, D.P., 2002a. A tenuous dust ring of Jupiter formed by escaping ejecta from the Galilean satellites. *J. Geophys. Res.* 107, E1. doi:10.1029/2000JE001434.
- Krivov, A.V., Wardinski, I., Spahn, F., Krüger, H., Grün, E., 2002b. Dust on the outskirts of the Jovian system. *Icarus* 157, 436–455.
- Krüger, H., 2003. *Jupiter's Dust Disc*, An Astrophysical Laboratory. Shaker Verlag Aachen, ISBN 3-8322-2224-3. Habilitation Thesis, Ruprecht-Karls-Universität, Heidelberg.
- Krüger, H., Bindschadler, D., Dermott, S.F., Graps, A.L., Grün, E., Gustafson, B.A., Hamilton, D.P., Hanner, M.S., Horányi, M., Kissel, J., Lindblad, B., Linkert, D., Linkert, G., Mann, I., McDonnell, J.A.M., Moissl, R., Morfill, G.E., Polansky, C., Schwehm, G.H., Srama, R., Zook, H.A., 2006. Galileo dust data from the jovian system: 1997 to 1999. *Planet. Space Sci.* 54, 879–910.
- Krüger, H., Grün, E., Graps, A.L., Bindschadler, D.L., Dermott, S.F., Fechtig, H., Gustafson, B.A., Hamilton, D.P., Hanner, M.S., Horányi, M., Kissel, J., Lindblad, B., Linkert, D., Linkert, G., Mann, I., McDonnell, J.A.M., Morfill, G.E., Polansky, C., Schwehm, G.H., Srama, R., Zook, H.A., 2001. One year of Galileo dust data from the jovian system: 1996. *Planet. Space Sci.* 49, 1285–1301.
- Krüger, H., Grün, E., Hamilton, D.P., Baguhl, M., Dermott, S.F., Fechtig, H., Gustafson, B.A., Hanner, M.S., Horányi, M., Kissel, J., Lindblad, B.A., Linkert, D., Linkert, G., Mann, I., McDonnell, J.A.M., Morfill, G.E., Polansky, C., Riemann, R., Schwehm, G.H., Srama, R., Zook, H.A., 1999a. Three years of Galileo dust data: II. 1993 to 1995. *Planet. Space Sci.* 47, 85–106.
- Krüger, H., Grün, E., Heck, A., Lammers, S., 1999b. Analysis of the sensor characteristics of the Galileo dust detector with collimated Jovian dust stream particles. *Planet. Space Sci.* 47, 1015–1028.
- Krüger, H., Grün, E., Linkert, D., Linkert, G., Moissl, R., 2005. Galileo long-term dust monitoring in the jovian magnetosphere. *Planet. Space Sci.* 53, 1109–1120.
- Krüger, H., Krivov, A.V., Grün, E., 2000. A dust cloud of Ganymede maintained by hypervelocity impacts of interplanetary micrometeoroids. *Planet. Space Sci.* 48, 1457–1471.
- Krüger, H., Krivov, A.V., Hamilton, D.P., Grün, E., 1999c. Detection of an impact-generated dust cloud around Ganymede. *Nature* 399, 558–560.
- Krüger, H., Krivov, A.V., Sremčević, M., Grün, E., 2003. Galileo measurements of impact-generated dust clouds surrounding the Galilean satellites. *Icarus* 164, 170–187.

- 1139 Landgraf, M., Baggeley, W.J., Grün, E., Krüger, H., Linkert, G., 2000. Aspects of the  
1140 mass distribution of interstellar dust grains in the Solar System from in situ  
1141 measurements. *J. Geophys. Res.* 105 (A5), 10343–10352. 1158
- 1142 Moissl, R., 2005. Galileos Staubmessungen in Jupiters Gossamer-Ringen. Diplom  
1143 thesis, Ruprecht-Karls-Universität, Heidelberg. 1159
- 1144 Ockert-Bell, M.E., Burns, J.A., Daubar, I.J., Thomas, P.C., Veverka, J., Belton, M.J.S.,  
1145 Klaasen, K.P., 1999. The structure of Jupiter's ring system as revealed by the  
1146 Galileo imaging experiment. *Icarus* 138, 188–213. 1160
- 1147 Showalter, M.R., Burns, J.A., Cuzzi, J.N., Pollack, J.B., 1985. Discovery of Jupiter's  
1148 'gossamer' ring. *Nature* 316, 526–528. 1161
- 1149 Showalter, M.R., de Pater, I., Verbanac, G., Hamilton, D.P., Burns, J.A., 2008.  
1150 Properties and dynamics of Jupiter's Gossamer rings from Galileo, Voyager,  
1151 Hubble and Keck images. *Icarus* 195, 361–377. 1162
- 1152 Srama, R., Kempf, S., Moragas-Klostermeyer, G., Helfert, S., Ahrens, T.J., Altobelli,  
1153 N., Auer, S., Beckmann, U., Bradley, J.G., Burton, M., Dikarev, V.V., Economou,  
1154 T., Fechtig, H., Green, S.F., Grande, M., Havnes, O., Hillier, J.K., Horanyi, M.,  
1155 Igenbergs, E., Jessberger, E.K., Johnson, T.V., Krüger, H., Matt, G., McBride, N.,  
1156 Mockler, A., Lamy, P., Linkert, D., Linkert, G., Lura, F., McDonnell, J.A.M.,  
1157 Möhlmann, D., Morfill, G.E., Postberg, F., Roy, M., Schwehm, G.H., Spahn, F.,  
1158 Svestka, Tschernjawski, V., Tuzzolino, A.J., Wäsch, R., Grün, E., 2006. In situ  
1159 dust measurements in the inner Saturnian system. *Planet. Space Sci.* 54, 967–  
1160 987. 1161
- 1161 Thiessenhusen, K.-U., Krüger, H., Spahn, F., Grün, E., 2000. Dust grains around  
1162 Jupiter—The observations of the Galileo Dust Detector. *Icarus* 144, 89–98. 1163
- 1163 Throop, H.B., Porco, C.C., West, R.A., Burns, J.A., Showalter, M.R., Nicholson, P.D.,  
1164 2004. The jovian rings: New results derived from Cassini, Galileo, Voyager and  
1165 Earth-based observations. *Icarus* 172, 59–77. 1166
- 1166 Williams, D.J., McEntire, R.W., Jaskulek, S., Wilken, B., 1992. The Galileo energetic  
1167 particles detector. *Space Sci. Rev.* 60, 385–412. 1168
- 1168 Willis, M.J., Burchell, M., Ahrens, T.J., Krüger, H., Grün, E., 2005. Decreased values of  
1169 cosmic dust number density estimates in the solar system. *Icarus* 176, 440–452. 1170
- 1170 Willis, M.J., Burchell, M., Cole, M., McDonnell, J.A.M., 2004. Influence of impact  
1171 ionization detection methods on determination of dust particle flux in space.  
1172 *Planet. Space Sci.* 52, 711–725. 1173
- 1173 Zeehandelaar, D.B., Hamilton, D.P., 2007. A local source for the Pioneer 10 and 11  
1174 Circumjovian dust detections, in: Krüger, H., Graps, A.L. (Eds.), *Dust in Planetary  
1175 Systems, Workshop, September 26–30, 2005, SP-643*. ESA Publications,  
1176 European Space Agency, Kauai, Hawaii, pp. 103–106. 1177