TODAY

- LAWS OF MOTION
- CONSERVATION LAWS
- **GRAVITY**

Extra credit (2 points)

- List Kepler's laws of planetary motion
- Be sure to include your name and section number
- You may consult your notes, but do not communicate with anyone else

Terms of Motion

Motion notions:



• Speed: Rate at which object moves speed = $\frac{\text{distance}}{\text{time}}$ (units of $\frac{\text{m}}{\text{s}}$)

example: speed of 10 m/s

- Velocity: Speed and <u>direction</u> example: 10 m/s, due east
- Acceleration: Rate of change in velocity

3

acceleration =
$$\frac{\text{speed}}{\text{time}}$$
 (units of $\frac{\text{m}}{\text{s}^2}$)

Momentum and Force

- Momentum = mass × velocity.
- A **net force** changes momentum, which generally means an acceleration (change in velocity).
- The rotational momentum of a spinning or orbiting object is known as **angular momentum**.

How is mass different from weight?

- Mass—the amount of matter in an object
- Weight—the *force* that acts upon an object



You are weightless in free-fall!

© 2007 Pearson Education Inc., publishing as Pearson Addison-Wesley

Why are astronauts weightless in space?



- There *is* gravity in space.
- Weightlessness is due to a constant state of free-fall.

6

© 2007 Pearson Education Inc., publishing as Pearson Addison-Wesley

What have we learned?

- How do we describe motion?
 - Speed = distance / time
 - Speed and direction => velocity
 - Change in velocity => acceleration
 - **Momentum** = mass × velocity
 - Force causes change in momentum, producing acceleration

What have we learned?

- How is mass different from weight?
 - —Mass = quantity of matter
 - —Weight = force acting on mass
 - —Objects are weightless in free-fall

4.2 Newton's Laws of Motion

Our goals for learning:

- How did Newton change our view of the universe?
- What are Newton's three laws of motion?

How did Newton change our view of the universe?



Sir Isaac Newton (1642–1727)

• He realized the same physical laws that operate on Earth also operate in the heavens:

 \Rightarrow one *universe*

- He discovered laws of motion and gravity.
- Much more: Experiments with light; first reflecting telescope, calculus...

Newton's three laws of motion



Newton's first law of motion: An object moves at constant velocity unless a net force acts to change its speed or direction. "Law of inertia"

In the absence of an applied force, an object at rest remains at rest. An object in motion remains in motion with a constant velocity.

Waiver Agreement

I understand that this demo is highly unlikely to result in agonizing pain for me, nor is it at all probable that the bones of my hand will be crushed to a fine powder, and my ligaments and tendons are almost certain to be okay as well. On the off chance that something does go horribly wrong, I agree to not hold the University of Maryland or, particularly, Professor Miller liable for any damages. Using the Force Which of the following motions does **not** require a net force? A.A car speeding up

B. A car moving at a constant speed in a circle

C. A car slowing down

D.A car moving at a constant speed in a straight line

E.I don't know

Newton's second law of motion

Force = mass × acceleration

F = ma

A force must be applied to change an object's state of motion (its momentum).

14

Newton's third law of motion

For every action, there is an equal and opposite reaction.



As the rocket fires, the shuttle doesn't just sit there. It accelerates Whenesetted, toothe wpjghtdiforce. balanced by the reaction force of the chair.



- Newton's Three Laws of Motion
 - 1. An object moves at constant velocity if no net force is acting.
 - 2. Force = mass \times acceleration.
 - 3. For every force, there is an equal and opposite reaction force.



Conserved Quantities

You can not destroy conserved quantities, only transfer them from one object to another.

- Mass
- Energy
- Momentum
- Angular momentum

Conservation of Momentum



- The total momentum of interacting objects cannot change unless an external force is acting on them.
- Interacting objects exchange momentum through equal and opposite forces.

What keeps a planet rotating and orbiting the Sun?



© 2006 Pearson Education, Inc., publishing as Addison Wesley

Conservation of Angular Momentum

angular momentum = mass × velocity × radius

L = mvr

- The angular momentum of an object cannot change unless an external twisting force (torque) is acting on it.
- Earth experiences no twisting force as it orbits the Sun, so its rotation and orbit will continue indefinitely.

Angular momentum conservation also explains why objects rotate faster as they shrink in radius:



e.g, kinetic energy:

• Energy makes matter move. $E_K = \frac{1}{2}mv^2$

• Energy is conserved, but it can... —transfer from one object to another. —change in form.

Basic Types of Energy

- Kinetic (motion)
- Radiative (light)
- Stored or potential

Energy can change type but cannot be destroyed.



Gravitational Potential Energy

- On Earth, it depends on...
 - an object's mass (m).
 - the strength of gravity (g).
 - the distance an object could potentially fall.



 $E_P = mgh$

Thermal Energy:

the collective kinetic energy of many particles (for example, in a rock, in air, in water)

Thermal energy is related to temperature but it is NOT the same.

Temperature is proportional to the *average* kinetic energy of the many particles in a substance.



higher temperature



Temperature Scales



Thermal energy is a measure of the total kinetic energy of all the particles in a substance. It therefore depends on both temperature AND density. Example:







Mass-Energy

Mass itself is a form of potential energy.

 $E = mc^2$

- A small amount of mass can release a great deal of energy.
- Concentrated energy can spontaneously turn into particles (for example, in particle accelerators).

Conservation of Energy

- Energy can be neither created nor destroyed.
- It can change form or be exchanged between objects.
- The total energy content of the universe is the same today as it was at the beginning of time.

Acceleration of Gravity

- All falling objects accelerate at the same rate (neglecting air resistance).
- On Earth, $g \approx 10 \text{ m/s}^2$: speed increases 10 m/s with each second of falling.

