

The Basic Structure of Present-Day Galaxies

1) Basic description of galaxy 'components'

- Stellar distribution: bulge, disk, bars,
- Distribution of gas (and dust)
- Dark matter halo

2) Parameter Relations in Galaxies

- Tully-Fisher, the 'Fundamental Plane' and the Kormendy relations
- Morphology, mass vs. kinematics
- Stellar mass vs. halo mass

3) Morphology and structure vs. formation history

- the sizes of disk galaxies
- the shapes of massive galaxies



4) Extreme ends of the galaxy property spectrum

- the smallest galaxies
- the most massive galaxies and galaxy clusters

5) Some comments on the Milky Way

(see IMPRS 2007 lecture notes on the web; UID/Pssw imprs-mwg/mwg-imprs)


<http://adc.gsfc.nasa.gov/mw>




Multiwavelength Milky Way




GALEX FUV + NUV (1500/2500 Å)



H α + R




IRAC 8.0 μ m



MIPS 24 μ m

Visible + Infrared



Sombrero Galaxy/Messier 104


NASA / JPL-Caltech / R. Kennicutt [University of Arizona], and the SINGS Team

Spitzer Space Telescope • IRAC

Visible: Hubble Space Telescope/Hubble Heritage Team


ssc2005-11a

M87 with HST




008

Interacting/merging galaxies



, 2008



Basic Description of the Stellar Distribution

- For fairly massive galaxies a basic two-component description of the stellar distribution proves useful:
 - **Bulges/spheroids**
 - **Disks**
- Radial profile description
Sersic (1968) profile

$$\Sigma(r) = \Sigma_e e^{-\kappa[(r/r_e)^{1/n} - 1]}$$
$$\kappa \approx 2n - 0.331$$



Disks: $n \sim 1$: 'exponential profile'

Spheroids: $n \sim 2-5$ ($n=4$: deVaucouleur)

NB: $n_{\text{spheroid}} = f(L_{\text{spheroid}})$

Note: bulge/disk approach (3D shape \leftrightarrow profile) not sensible for low-mass galaxies

2MASS large galaxy atlas



Radial profiles: Comments

- Many (massive) ellipticals fit the de Vaucouleur's profile beautifully
- Bulge-disk decompositions on the basis of radial profiles alone are terribly prone to fitting degeneracies





Figure 2: Decomposition of NGC2708 with both models. (a) (Left panel): $r^{1/4}$ bulge. (b) (Right panel): Exponential bulge.

Bulge-to-Disk Ratios in the Present-Day Universe (e.g. Tasca and White 2005 and Benson et al 2007 based on SDSS)

- Bulge-disk decomposition is interesting, because
 - a) disk-stars: no violent relax.
 - b) Spheroid stars: post-violent relax
 - c) $M_{\text{BH}} \sim M_{\text{Spheroid}}$



Globally:

60% of r-band light from disks
40% from spheroids

In stellar mass:



40%/60% disk/spheroid mass fraction

Spheroids dominate the massive end



Structure of Galaxy Disks I



- Vertical stellar profile can often be described by $f(z) = \text{sech}^{2/N}(Nz/z_0)$,
- In most galaxy disks a description by two vertical components is suggested (incl. Milky Way)
 - Thicker \rightarrow more (vertical) kinetic energy \rightarrow Why?
- (Some) stellar disks are ‘truncated’ in radius
 - Max. angular momentum, or
 - Threshold in star-formation efficiency?



Structure of Galaxy Disks II

- Stellar bars are common in disk galaxies
 - Often only recognized in near-IR images (less dust)
 - Consequence of disk instability
 - Effective means of angular momentum transport
 - Spiral arms are common and coherent features
 - even after accounting for young stars
- e.g. M51, Rix and Rieke 1995

SDSS (NYU Catalog)



The 3D Shapes of Spheroidal Galaxies

- What is the relation between intrinsic shape and projected ellipticity


axially symmetric case (oblate or prolate, see Binney/Merrifield):

$$q_{\text{internal}}^2 \sin^2 i + \cos^2 i = \begin{cases} q_{\text{projected}}^2 & (\text{oblate}) \\ 1/q_{\text{projected}}^2 & (\text{prolate}) \end{cases}$$

- If we view a sample from random angles, then $\cos(i)$ is uniform →

$$\underline{a : b : c \approx 1 : 0.95 : 0.7}$$



- Massive spheroidal galaxies are nearly oblate and only somewhat flat



Color-gradients in Galaxies


- Almost all galaxies become bluer outward
- Combination of
 - Decreasing dust
 - Decreasing age(?)
 - Decreasing metallicity
- Spheroids are redder/older than disks

Ensemble of spiral galaxies with dust and red bulges at the center →
(de Jong 1996)



Line-strength gradient (Mg2)
Davies et al 1993



(dust-free)
massive
spheroids



Basic Kinematics of Spheroidal Galaxies

Stellar velocity fields for nearby spheroidal galaxies

(Capellari, deZeeuw, Bacon, et al 2005)



- Generically:
 - Rotation rises slowly outwards
 - Dispersion falls gently outward
- $0 < v/\sigma < 1.5$

„Interstellar Gas“ in Galaxies

Interstellar gas occurs in a wide range of physical conditions

Name	State	Density	Temperature	Main diagnostics
„hot“	fully ionized	10^{-2} cm^{-3}	10^6 K	X-rays UV absorption
warm (H II)	fully ionized	1 cm^{-3}	10^4 K	Optical emission lines
neutral (H I)	neutral atomic	1 cm^{-3}	10^2 K	21cm line
„cold“ molecular gas	molecular	100	20 K	CO lines (radio, sub-mm)


What sets the temperature and the physical state of the gas?

Heating processes

- photo-ionization
- mechanical (shock) heating

Cooling processes

- Bremsstrahlung
- line cooling




HWR April 1, 2008

Galaxies and their Dark Matter Halos

- All evidence for dark matter halos on galaxy scales comes from the comparison/modeling of kinematic tracers with identified mass components.
- *Kinematic tracers:* stars, cold gas (HI and H α), hot gas (X-ray), satellites (GCs and galaxies) and photons (gravitational lensing)
- *Identified (baryonic) mass components:* stars, hot gas (in clusters), cold gas (\sim 10% of stars)
- *Historically:*
 - Need for dark matter from dynamics on scales of galaxies played an enormous role in establishing its (dynamical) existence
- *Current Paradigm:*
 - Dark matter is a indispensable ingredient in structure formation; galaxies are the places where DM is least dominant → DM studies on galaxy scales can be tricky



Observational Constraints on Dark Matter Halos around Big Galaxies

a) HI rotation curves
„flat“ to ~ 30 kpc ($\gg R_{\text{opt}}$)




$$V_c^2(R) = -R \frac{d \Phi(\vec{r})}{d R} \\ \approx -g M(< R) / R$$

Model fits to HI rotation curves




Note:
can't constrain 2 functions, $\rho_*(r)$ and $\rho_{DM}(r)$,
by 1 function $V_c(r)$



Rotation Curves are “Flat”

- At small radii the stars (probably) dominate
- At large radii the dark matter dominates
- No obvious reason at first that the rotation curve does not show the transition from the star to the DM dominated regime



Dark Matter Halos at large radii



Use satellite galaxies (Zaritsky 1994)

e.g. Prada et al 2002 by stacking images

→ DM halos around MW-like galaxies extend to >200kpc

NFW profile:

$$\rho(r) = \delta_s / [(r/r_s) (1 + r/r_s)^2]$$




Present-Day knowledge about Dark Matter Halos

c) Strong and weak gravitational lensing

(Maoz and Rix 1993; Brainerd et al 1988;
McKay et al 2001)


Background galaxy lensed into
arcs by lens



(Sheldon et al.
2004; SDSS)

Correlation function between

- a) Shear (from gravitational lensing seen in background images)
- b) Position of foreground galaxies




X-ray gas around massive galaxies

- Only in massive galaxies (and galaxy clusters) is the 'hot' phase hot enough to be detected by current X-ray satellites
- Assume gas is in (approximate) hydrostatic equilibrium

$$\ln\left(\frac{\rho_g}{\rho_{g0}}\right) = -\ln\left(\frac{T}{T_0}\right) - G\mu m_p \int_{R_0}^R \frac{M_{grav}(< R)}{kTR^2} dR$$

- In all massive galaxies with good measurements:
DM halo with properties expected from Λ CDM (NFW halo)



2) 'Parameter Relations' in (Present-Day) Galaxies


Many parameters with which to describe the stellar component of galaxies are tightly correlated
(though such correlations are/were not 'expected')

Most of them can be cast as

(stellar) luminosity/mass vs

- size
- characteristic velocity (Tully-Fisher; Faber-Jackson)
- 3D - shape
- (radial) concentration, black hole mass



These correlations are important constraints on galaxy formation mechanisms



John Kormendy has been a pioneer in pointing out
that the photometric descriptions are correlated

The ‘Tully-Fisher’ Relation for Disk Galaxies

- Tully&Fisher 1977
 - HI linewidth correlates well with absolute magnitude of spiral galaxy.
- In general:
 - Correlation between circular velocity and stellar luminosity
 - L_{opt} can predict v_{circ} to $\sim 5\text{-}8\%$
 - $M_*, L_{\text{opt}} \sim v_c^{3\text{-}4}$
- Historically: extremely important distance indicator
- Now: also constraint on galaxy formation



Stellar Mass (SED + Kroupa IMF)

Explanations for a Tully-Fisher-like relation

- Let's consider the self-gravitating case

$$v_{\max}^2 \sim G\Sigma_0 R_d \quad \text{with } \Sigma_0 \text{ central mass density and } R_d \text{ scale length}$$

$$\Rightarrow L \sim v_{\max}^4 I_0^{-1} \Gamma_{disk}^{-2} \quad \text{with } I_0 = \Sigma_0 / \Gamma_{disk}$$

Right slope, but central surface brightness/mass density should be a 3rd parameter

- Let's presume the disk is a small fraction assembled from a DM halo
For the halo (also Mo, Mao and White 1993)

$$r_{\text{vir}} = \sqrt{\frac{2}{\Delta_{\text{vir}}(z)}} \frac{V_{\text{vir}}}{H(z)} \quad \text{and} \quad M_{\text{vir}} = \sqrt{\frac{2}{\Delta_{\text{vir}}(z)}} \frac{V_{\text{vir}}^3}{G H(z)} \quad \rho_{\text{crit}} = 3H^2(z)/(8\pi G)$$


$$\Delta_{\text{vir}}(z)$$

Factor by which density within virial radius exceeds mean density of the universe

$$M_d \approx 1.3 \times 10^{11} h^{-1} M_\odot \left(\frac{m_d}{0.05} \right) \left(\frac{V_{\text{vir}}}{200 \text{ km s}^{-1}} \right)^3$$


with NO surface brightness/mass dependence!

Parameter relations for (massive) spheroids: Faber-Jackson and the ‘fundamental plane’




- For spheroids: →
3-parameter relation!!
- $M/L = f(M)$

One version of the ‘fundamental plane’, involving L, R_e, σ_*





Rotation support and isophote shape = $f(L)$



Stellar mass vs. Halo Mass


How efficient is galaxy formation?

Dark matter halos cluster ...
more massive \rightarrow more clustered



Analogous for galaxies

Identify (observed) galaxy populations with (simulated) halos that have the same clustering properties



M/L strong function of M!

Van den Bosch et al 2006
HWR April 1, 2008

3) Galaxy Structure vs Formation Mechanisms

a) (Disk) Galaxy Sizes and Angular Momentum

- In disk galaxies the stellar body is centrifugally supported (stars move on near-circular orbits), with a "spin parameter",

$$\lambda \equiv \frac{J}{G} \sqrt{\frac{E}{M^5}}$$

where J is the angular momentum and E is the binding energy of the system.

$\lambda_{*, \text{observed}} \approx 0.5 - 1$ for disks ($\lambda_* \approx 0.005$ for spheroids)

→ Disk size is given by the angular momentum of the material

Angular momentum comes from torques of (adjacent) mass distribution (Hoyle 1949, Ryden and Gunn 1987)

Linear theory: $\lambda_{\text{total}} = \lambda_{\text{DM}} = \lambda_{\text{gas}}(\text{init}) \sim 1/20$

Why galaxies are disks of a characteristic size

- Torques before the collapse induce spin $\lambda \sim 0.07$
 - **Gas dissipates (by radiation) all the energy it can without violating angular momentum conservation \rightarrow circular orbit**
 - Fall and Efstathion (1980) showed that observed galaxy disks ($\lambda \sim 0.5$) can form only in DM halos through dissipation
 \rightarrow central concentration (J conserved) \rightarrow spin-up.
- a) Presume there is no DM:

$$\lambda_{obs} = \frac{J \cdot E^{1/2}}{GM^{5/2}} = \lambda_{init} \sqrt{\frac{R_{init}}{R}} \quad \Rightarrow \quad \frac{R_{init}}{R} \approx 50$$

We observe $M_{disk} \approx 5 \times 10^{10} M_{sun}$, $R_{disk} \approx 8 \text{ kpc} \Rightarrow R_{init} \approx 400 \text{ kpc}$

$\Rightarrow R_{turn-around} \approx 2 R_{init} \approx 800 \text{ kpc}$

$\Rightarrow t_{collapse} \sim 50 \cdot 10^9 \text{ years}$ for $M \sim 5 \times 10^{10} M_{sun}$

b) If the gas is only a small fraction of the total mass:

$\Rightarrow v_c(r)$ remains unchanged

$$\Rightarrow R_{init}/R \sim \frac{\lambda_{obs}}{\lambda_{init}} \Rightarrow R_{init} \sim 80 \text{ kpc}$$

$$\Rightarrow t_{dyn} \sim 10^9 \text{ years}$$

and there is enough time to form disks.

It turns out that the assumption of angular momentum conservation during the gas dissipation yield disk sizes as observed (assuming $\lambda \sim 0.07$)

However: in (numerical) simulations much of the angular momentum is lost \rightarrow modelled disks too small (unsolved)

Luminosity/Mass vs. Size


[state of the art incarnation: Shen et al 2004 based on SDSS]

- Well defined size relations with ~ 2.5 scatter

$$R \sim M^{0.5}$$

at $M > 3 \times 10^{10} M_{\odot}$

- Galaxy (stellar) sizes are related to the characteristic angular momentum of the stars (see below)



Why are massive galaxies spheroids?

1. Stars form from dense, cold gas
 - either in disks
 - or from gas that is (violently) shock compressed
 2. In the established cosmological paradigm larger (halos) form from the coalescence of smaller units
- Stars in an (near) equilibrium system form from a disk and stay disk-like
- 'Violent relaxation' shaking up stars (or stars formed during such an event) end up in spheroids

Is it plausible that in nearly all massive galaxies a (major) merger occurred after star-formation was largely complete?





Fig. 4.—Evolution of the stellar distribution in encounter A, projected onto the orbital plane. The scale is the same as in Fig. 3.


Some physics of mergers



+ Some gas dissipation is needed to get the (central) densities of ellipticals 'right'

Merging moves objects 'within' the fundamental plane!

Isophote shapes Naab&Burkert simulations




The Smallest and the Largest Galaxies

- Questions:
 - Is there an empirical (upper/lower) limit to 'galaxies'?


1) 'Dwarf' Galaxies

- Definition (not universally established):
 - galaxy that has $< 1/10^{\text{th}}$ of L_* (or M_*) Milky_Way
 - or $v_{\text{circ}} < 100 \text{ km/s}$
- Most abundant type of galaxies; contributes negligibly to the total stellar mass budget.
- Structure and morphologies
 - Often 'irregular' (highly asymmetric)
 - Often of very low surface brightness
- Stellar populations
 - Inevitably low metallicity ($< 1/10$ solar)
 - Some have very young pops. (most stars after $z < 0.5$)
 - Some have only old ($> 10 \text{ Gyrs}$) stars, many are mixed populations
- Dwarfs are interesting regime to test gravity $\leftarrow \rightarrow$ 'feed-back'




The Faintest Galaxies Known to Date Milky Way Satellites

- Most found recently by SDSS (e.g. Belokurov et al 2006)
- Seem to be dark matter dominated
- Same luminosity as globular clusters, but 1000x lower stellar surface mass density



The Extreme Limit of “Galaxy Clustering” **Galaxy Clusters**



Galaxy Cluster Abell 2218

NASA, A. Fruchter and the ERO Team (STScI, ST-ECF) • STScI-PRC00-08



HST • WFPC2

April 1, 2008

cD galaxies: the most massive galaxies

- Galaxies with $\geq 5 L_{\text{MW}}$ are only found at the centers of galaxy clusters
- They are without exception spheroidal
- They have extended light profiles that blend into 'intra-cluster' light

'stacked' image of 700 cluster centers (Zibetti et al 2005)



Virgo Cluster:

- Nearest large galaxy cluster with more than 2000 galaxies brighter than $M_B \simeq -14$ ($L_B \sim 10^{7.8} L_\odot$)
- Distance $\sim 17 Mpc$ (dependent on H_0)
- Extend $\sim 10^\circ \doteq 3Mpc \times 3Mpc$
- Irregular cluster, densest regions dominated by ellipticals
- Velocity dispersion of galaxies about 600km/s


Coma Cluster:

- One of the most luminous clusters known
- Distance $\sim 100 Mpc$ (dependent on H_0)
- Regular cluster with probably subcluster merging from SW
- Dominated by ellipticals and S0s, two central cDs and one in subcluster
- Velocity dispersion of galaxies about 1000km/s
- Strong X-ray source

Virgo Cluster of Galaxies D \sim 15Mpc

HWR April 1, 2008

Galaxy Clusters are filled with Hot Gas



Coma cluster (left: optical image, right: X-ray image)

The X-ray spectra show the characteristics of Bremsstrahlung of a $\sim 10^8 K$ hot gas.

The volume emissivity is: $\frac{dP}{dV} = 2.410^{-27} T^{-1/2} N_e^2 \left[\frac{\text{erg}}{\text{cm}^3 \text{s}} \right]$

The cooling time of the plasma is: $t_{cool} = \frac{3N_e kT}{\frac{dP}{dV}} \simeq \frac{10^{11}}{N_e} T^{1/2} [\text{s}]$

For the center of the Coma cluster we have:

$$L \simeq 10^{44} \text{erg/s}$$

$$\overline{n_e} \simeq 10^{-3} \text{cm}^{-3}$$

$$\overline{\tau_{cool}} \simeq 10^{10} \text{yrs}$$

$$M_{gas} \simeq 10^{13} M_\odot$$

The following correlations exist between the different components of galaxy clusters:


- The central galaxy density is higher for higher L_X .
- The fraction of spirals is lower for higher L_X .
- The temperature T is proportional to L_X and typically 10^8K .
- The gas metallicity is lower for higher T and typically 1/3 of solar.
- The ratio of gas-mass to galaxy-mass increases with T up to 5 or more.
- The dominant component in all clusters is dark matter. This follows consistently from the dynamics of galaxies, the hydrostatic equilibrium of the X-ray gas and from gravitational lensing. The typical mass ratios are:

$$\text{galaxies : X-ray-gas : dark-matter} \simeq 1 : 5 : 25$$

Galaxy Properties: Summary


- Description of the stellar body of galaxies
 - Spheroid : post-violent relaxation/merger stars
 - Disks: not shook-up since formation
 - In the present-day universe ~50%/50
- The variety of galaxies in
 - Morphology, shape, structure, ...
 - Stellar content and metallicity....

is very restricted (many parameter relations).
- The stellar mass range of galaxies
 - Clearly limited at upper end: 'brightest cluster galaxies'
 - No end in sight (<1000 M*) at the lower end?
- Milky Way
 - Typical galaxies in many ways
 - Central in shaping our thinking about galaxies
 - (historically, but also in future)




HWR April 1, 2008

5. The Thick and Thin Disk





30.08.2007

E.K. Grebel: The M



HWR April 1, 2008

5. The Thick and Thin Disk



Age-metallicity evolution of the different Milky Way components (same color coding as in schematic above).

Buser 2000



29


30.08.2007

E.K. Grebel: The Milky Way and the Local Group (II)

HWR April 1, 2008

11. The Stellar Halo

Carney et al. (1996) survey of proper motion stars :
Orbital eccentricity vs. $[m/H]$ – modern version of ELS



Striking kinematical difference between disk and halo

31.08.2007

E.K. Grebel: The Milky Way and the Local Group (IV)



10

HWR April 1, 2008

7 Associations and Open Star Clusters



Photo: Gendler



Photo: Bessell

★ Open Star Clusters

- Mostly loose diffuse systems in the Galactic disk.
- Masses: $3 \leq \log M [M_{\odot}] \leq 5$, $\gg 100$ stars, $100-30,000 L_{\odot}$
- Radii: 1 to 20 pc; in Galactic disk, small scale height.
- Average life time: ~ 300 Myr; only weakly bound, internal dynamics dominate dissolution.

13. The Galactic Bulge



Milky Way:
Small **exponential** bulge –
typical of later-type galaxies.

(Unlike the large $r^{1/4}$ bulge of M31)




01.09.2007

E.K. Grebel: Th



The Local Group



HWR April 1, 2008